

A/Box  
Sep

Atty Dkt No. 5995.US.P1  
0200-0053.20

## APPLICATION TRANSMITTAL LETTER



Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

☒ "Express Mail" Mailing Label No. EM 049 296283 US, Date of Deposit OCTOBER 31, 1997  
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

PATRICIA K. HIMENES  
(Typed or Printed Name of Person Mailing Paper or Fee)

Patricia K. Himenes  
(Signature of Person Mailing Paper or Fee)

Transmitted herewith for filing is the patent application of Patricia A. BILLING-MEDEL, Maurice COHEN, Tracey L. COLPITTS, Paula N. FRIEDMAN, Julian GORDON, Edward N. GRANADOS, Steven C. HODGES, Michael R. KLASS, Jon D. KRATOCHVIL, Lisa ROBERTS-RAPP, John C. RUSSELL and Steven D. STROUPE for REAGENTS AND METHODS USEFUL FOR DETECTING DISEASES OF THE BREAST, which is a continuation-in-part of U.S. serial no. 08/742,067, filed October 31, 1996.

Enclosed are:

- 6 sheet(s) of X formal \_\_\_ informal drawing(s).
- \_\_\_ A claim for foreign priority under 35 U.S.C. § 119/363 in  
\_\_\_ a separate document \_\_\_ the declaration.
- \_\_\_ A claim for priority under 35 U.S.C. § 119(e)(1) in  
\_\_\_ a separate document \_\_\_ the declaration.
- \_\_\_ A certified copy of the priority document.
- \_\_\_ Verified Statement(s) Claiming Small Entity Status.
- X Other: Sequence Listing in paper form (pp. 89-95); Sequence Listing in computer readable form; and Statement to Support Filing and Submission in Accordance with 37 C.F.R. §§ 1.821-1.825.

The declaration of the inventor X is enclosed X unsigned.

The fee has been calculated as follows:

|  |           |           |
|--|-----------|-----------|
| A. Basic Application Fee   |           | \$790.00  |
| B. Total Claims 39 - 20 = 19   | x \$22.00 | 418.00    |
| C. Independent Claims 20 - 3 = 17  | x \$82.00 | 1394.00   |
| D. If multiple dependent claims present, add   | \$270.00  | 0.00      |
| E. Total Application Fee (Total of A, B, C, & D)   | =         | 2602.00   |
| <hr/>  |           |           |
| F. If verified statement of small entity status is enclosed, reduce Total Application Fee by 50% |           | NA        |
| G. Application Fee Due (E - F)   | =         | 2602.00   |
| <hr/>  |           |           |
| H. Assignment Recording Fee of \$40.00 if assignment document is enclosed                        | \$40.00   | NA        |
| <hr/>  |           |           |
| I. TOTAL FEE (G + H)   |           | \$2602.00 |

Respectfully submitted,

Date: 31 OCTOBER 1997

By: Thomas P. McCracken  
Thomas P. McCracken  
Registration No. 38,548  
Attorney for Applicants

ABBOTT LABORATORIES  
100 Abbott Park Road  
Abbott Park, IL 60064-3500  
Telephone: (847) 937-6100  
Facsimile: (847) 938-2623

Atty Dkt 5995.US.P1  
0200-0053.20  
PATENT

"Express Mail" Mailing Label No. EM 049 296 283 US  
Date of Deposit OCTOBER 21, 1997

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

PATRICIA K. HIMENES  
Typed or Printed Name of Person Mailing Paper or Fee

Patricia K. Himenes  
Signature of Person Mailing Paper or Fee

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

BILLING-MEDEL et al.

Serial No.: CIP of 08/742,067

Group Art Unit: Unassigned

Filing Date: even date herewith

Examiner: Unassigned

Title: REAGENTS AND METHODS USEFUL FOR DETECTING DISEASES  
OF THE BREAST

STATEMENT TO SUPPORT FILING AND SUBMISSION IN ACCORDANCE  
WITH 37 C.F.R. §§ 1.821-1.825

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

The undersigned hereby states that the content of the attached papers and the computer-readable copy of the Sequence Listing, submitted in accordance with 37 C.F.R. §§ 1.821(c) and (e), respectively, are the same.

Respectfully submitted,

Date: 31 OCTOBER 1997

By: Thomas P. McCracken  
Thomas P. McCracken  
Registration No. 38,548  
Attorney for Applicants

ABBOTT LABORATORIES  
100 Abbott Park Road  
Abbott Park, IL 60064-3500  
Telephone: (847) 937-0378  
Facsimile: (847) 938-2623

00962004 103 46029680

CERTIFICATE OF MAILING BY "FIRST CLASS MAIL"  
"First Class Mail" and by "FIRST CLASS MAIL"  
Date of Deposit: OCTOBER 31 1997  
I hereby certify that this paper or the mailing document  
with the United States Postal Service "First Class Mail Post  
Office to Addressee" covers under 37 U.S.C. 1111 on the date  
indicated above and is addressed to the Assistant  
Comptroller for Records, Washington, D.C. 20531

Atty Dkt No.5995.US.P1  
PATENT

-1-

PATRICIA K. HIMENES  
(Typed or Printed Name of Person Mailing Paper or Fee)  
Patricia K. Himenes  
(Signature of Person Mailing Paper or Fee)

5

REAGENTS AND METHODS USEFUL FOR  
DETECTING DISEASES OF THE BREAST

10 Cross-Reference to Related Application

This application is a continuation-in-part of U.S. application Serial No.  
08/742,067, filed October 31, 1996, from which priority is claimed pursuant to 35  
U.S.C. §120 and which is incorporated herein by reference in its entirety.

15 Background of the Invention

The invention relates generally to detecting diseases of the breast, and  
more particularly, relates to reagents such as polynucleotide sequences and the  
polypeptide sequences encoded thereby, as well as methods which utilize these  
sequences, which are useful for detecting, diagnosing, staging, monitoring,  
20 prognosticating, preventing or treating, or determining predisposition to diseases  
or conditions of the breast such as breast cancer.

Breast cancer is the most common form of cancer occurring in females in  
the US. The incidence of breast cancers in the United States is projected to be  
180,200 cases diagnosed and 43,900 breast cancer related deaths to occur during  
25 1997 (American Cancer Society statistics). Worldwide, the incidence of breast  
cancer has increased from 700,000 in 1985 to about 900,000 in 1990. G.N.  
Hortobagyi et al., CA Cancer J Clin 45: 199-226 (1995).

Procedures used for detecting, diagnosing, staging, monitoring,  
prognosticating, preventing or treating, or determining predisposition to diseases  
30 or conditions of the breast such as breast cancer are of critical importance to the  
outcome of the patient. For example, patients diagnosed with early breast cancer  
have greater than a 90% five-year relative survival rate as compared to a survival  
rate of about 20% for patients diagnosed with distantly metastasized breast  
cancers. (American Cancer Society statistics). Currently, the best initial  
35 indicators of early breast cancer are physical examination of the breast and

08962094-103197

46 PATENT "16029680

mammography. J.R. Harris et al. In: Cancer: Principles and Practice of Oncology, Fourth Edition, pp. 1264-1332, Philadelphia, PA: J/B. Lippincott Co. (1993). Mammography may detect a breast tumor before it can be detected by physical examination, but it has limitations. For example, mammography's predictive value depends on the observer's skill and the quality of the mammogram. In addition, 80 to 93% of suspicious mammograms are false positives, and 10 to 15% of women with breast cancer have false negative mammograms. C.J. Wright et al., Lancet 346: 29-32 (1995). New diagnostic methods which are more sensitive and specific for detecting early breast cancer are clearly needed.

Breast cancer patients are closely monitored following initial therapy and during adjuvant therapy to determine response to therapy, and to detect persistent or recurrent disease, or early distant metastasis. Current diagnostic procedures for monitoring breast cancer include mammography, bone scan, chest radiographs, liver function tests and tests for serum markers. The serum tumor markers most commonly used for monitoring patients are carcinoembryonic antigen (CEA) and CA 15-3. Limitations of CEA include absence of elevated serum levels in about 40% of women with metastatic disease. In addition, CEA elevation during adjuvant therapy may not be related to recurrence but to other factors that are not clinically important. CA 15-3 can also be negative in a significant number of patients with progressive disease and, therefore, fails to predict metastasis. Both CEA and CA 15-3 can be elevated in nonmalignant, benign conditions giving rise to false positive results. Therefore, it would be clinically beneficial to find a breast associated marker which is more sensitive and specific in detecting cancer recurrence. J. R. Harris, et al., supra. M. K. Schwartz, In: Cancer: Principles and Practice of Oncology, Vol. 1, Fourth Edition, pp. 531 - 542, Philadelphia, PA: J/B. Lippincott Co. 1993.

Another important step in managing breast cancer is to determine the stage of the patient's disease, because it has potential prognostic value and provides criteria for designing optimal therapy. Currently, pathological staging of breast cancer is preferable over clinical staging because the former gives a more accurate prognosis. J. R. Harris, et al., supra. On the other hand, clinical staging would be preferred were it at least as accurate as pathological staging, because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of breast cancer could be improved by detecting

new markers in serum or urine which could differentiate between different stages of invasion. Such markers could be mRNA or protein markers expressed by cells originating from the primary tumor in the breast but residing in blood, bone marrow or lymph nodes and could serve as sensitive indicators for metastasis to these distal organs. For example, specific protein antigens and mRNA, associated with breast epithelial cells, have been detected by immunohistochemical techniques and RT-PCR, respectively, in bone marrow, lymph nodes and blood of breast cancer patients suggesting metastasis. K. Pantel, et al., Onkologie 18: 394-401 (1995).

Such procedures also could include assays based upon the appearance of various disease markers in test samples such as blood, plasma, serum, or urine obtained by minimally invasive procedures which are detectable by immunological methods. These procedures would provide information to aid the physician in managing the patient with disease of the breast at low cost to the patient. Markers such as prostate specific antigen (PSA) and human chorionic gonadotropin (hCG) exist and are used clinically for screening patients for prostate cancer and testicular cancer, respectively. For example, PSA normally is secreted by the prostate at high levels into the seminal fluid, but is present in very low levels in the blood of men with normal prostates. Elevated levels of PSA protein in serum are used in the early detection of prostate cancer or disease in asymptomatic men. See, for example, G.E. Hanks, et al., In: *Cancer: Principles and Practice of Oncology*, Vol. 1, Fourth Edition, pp. 1073-1113, Philadelphia, PA: J.B. Lippincott Co. 1993. M. K. Schwartz, et al., In: *Cancer: Principles and Practice of Oncology*, Vol. 1, Fourth Edition, pp. 531-542, Philadelphia, PA: J.B. Lippincott Co. 1993. Likewise, the management of breast diseases could be improved by the use of new markers normally expressed in the breast but found in elevated amounts in an inappropriate body compartment as a result of the disease of the breast.

Further, new markers which could predict the biologic behavior of early breast cancers would also be of significant value. Early breast cancers that threaten or will threaten the life of the patient are more clinically important than those that do not or will not be a threat. G.E. Hanks, supra. Such markers are needed to predict which patients with histologically negative lymph nodes will experience recurrence of cancer and also to predict which cases of ductal carcinoma in situ will develop into invasive breast carcinoma. More accurate prognostic markers would allow the clinician to accurately identify early cancers

localized to the breast which will progress and metastasize if not treated aggressively. Additionally, the absence of a marker for an aggressive cancer in the patient could spare the patient expensive and non-beneficial treatment. J. R. Harris, et al., supra. E. R. Frykberg, et al., Cancer 74: 350-361 (1994).

5           It would be advantageous, therefore, to provide specific methods and reagents useful for detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining predisposition to diseases or conditions of the breast. Such methods would include assaying a test sample for products of a gene which are overexpressed in diseases and conditions associated with the  
10   breast including cancer. Such methods may also include assaying a test sample for products of a gene which have been altered by the disease or condition associated with the breast including cancer. Such methods may further include assaying a test sample for products of a gene whose distribution among the various tissues and compartments of the body have been altered by a breast-  
15   associated disease or condition including cancer. Such methods would comprise making cDNA from mRNA in the test sample, amplifying, when necessary, portions of the cDNA corresponding to the gene or a fragment thereof, and detecting the cDNA product as an indication of the presence of the disease or condition including cancer or detecting translation products of the mRNAs  
20   comprising gene sequences as an indication of the presence of the disease. Useful reagents include polynucleotides, or fragments thereof which may be used in diagnostic methods such as reverse transcriptase-polymerase chain reaction (RT-PCR), PCR, or hybridization assays of mRNA extracted from biopsied tissue, blood or other test samples; or proteins which are the translation  
25   products of such mRNAs; or antibodies directed against these proteins. Such assays would include methods for assaying a sample for products of the gene and detecting the products as an indication of disease of the breast. Drug treatment or gene therapy for diseases and conditions of the breast including cancer can be based on these identified gene sequences or their expressed  
30   proteins, and efficacy of any particular therapy can be monitored. Furthermore, it would be advantageous to have available alternative, non-surgical diagnostic methods capable of detecting early stage breast disease such as cancer.

Summary of the Invention

The present invention provides a method of detecting a target BS106 polynucleotide in a test sample which comprises contacting the test sample with at least one BS106-specific polynucleotide and detecting the presence of the target  
5 BS106 polynucleotide in the test sample. The BS106-specific polynucleotide has at least 50% identity with a polynucleotide selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. Also, the BS106-specific polynucleotide may be attached to a solid phase prior to performing  
10 the method.

The present invention also provides a method for detecting BS106 mRNA in a test sample, which comprises performing reverse transcription (RT) with at least one primer in order to produce cDNA, amplifying the cDNA so obtained using BS106 oligonucleotides as sense and antisense primers to obtain BS106 amplicon,  
15 and detecting the presence of the BS106 amplicon as an indication of the presence of BS106 mRNA in the test sample, wherein the BS106 oligonucleotides have at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. Amplification can be  
20 performed by the polymerase chain reaction. Also, the test sample can be reacted with a solid phase prior to performing the method, prior to amplification or prior to detection. This reaction can be a direct or an indirect reaction. Further, the detection step can comprise utilizing a detectable label capable of generating a measurable signal. The detectable label can be attached to a solid phase.

25 The present invention further provides a method of detecting a target BS106 polynucleotide in a test sample suspected of containing target BS106 polynucleotides, which comprises (a) contacting the test sample with at least one BS106 oligonucleotide as a sense primer and at least one BS106 oligonucleotide as an anti-sense primer, and amplifying same to obtain a first stage reaction product; (b)  
30 contacting the first stage reaction product with at least one other BS106 oligonucleotide to obtain a second stage reaction product, with the proviso that the other BS106 oligonucleotide is located 3' to the BS106 oligonucleotides utilized in step (a) and is complementary to the first stage reaction product; and (c) detecting the second stage reaction product as an indication of the presence of a target BS106  
35 polynucleotide in the test sample. The BS106 oligonucleotides selected as reagents



in the method have at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. Amplification may be performed by the polymerase chain reaction. The test sample can be reacted either directly or indirectly with a solid phase prior to performing the method, or prior to amplification, or prior to detection. The detection step also comprises utilizing a detectable label capable of generating a measurable signal; further, the detectable label can be attached to a solid phase. Test kits useful for detecting target BS106 polynucleotides in a test sample are also provided which comprise a container containing at least one BS106-specific polynucleotide selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. These test kits further comprise containers with tools useful for collecting test samples (such as, for example, blood, urine, saliva and stool). Such tools include lancets and absorbent paper or cloth for collecting and stabilizing blood; swabs for collecting and stabilizing saliva; and cups for collecting and stabilizing urine or stool samples. Collection materials, such as papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. The collection materials also may be treated with or contain preservatives, stabilizers or antimicrobial agents to help maintain the integrity of the specimens.

The present invention provides a purified polynucleotide or fragment thereof derived from a BS106 gene. The purified polynucleotide is capable of selectively hybridizing to the nucleic acid of the BS106 gene, or a complement thereof. The polynucleotide has at least 50% identity to a polynucleotide selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. Further, the purified polynucleotide can be produced by recombinant and/or synthetic techniques. The purified recombinant polynucleotide can be contained within a recombinant vector. The invention further comprises a host cell transfected with said vector.

The present invention further provides a recombinant expression system comprising a nucleic acid sequence that includes an open reading frame derived from BS106. The nucleic acid sequence has at least 50% identity with a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2,

SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. The nucleic acid sequence is operably linked to a control sequence compatible with a desired host. Also provided is a cell transfected with this recombinant expression system.

5           The present invention also provides polypeptides encoded by BS106. The polypeptides can be produced by recombinant technology, provided in purified form, or produced by synthetic techniques. The polypeptides comprise amino acid sequences which have at least 50% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments of SEQUENCE ID NOS 17-20.

Also provided is an antibody which specifically binds to at least one BS106 epitope. The antibody can be a polyclonal or monoclonal antibody. The epitope is derived from an amino acid sequence selected from the group consisting of

SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof. Assay kits for determining the presence of BS106 antigen or anti-BS106 antibody in a test sample are also included. In one embodiment, the assay kits comprise a container containing at least one BS106 polypeptide having at least 50% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof. Further, the test kit can comprise a container with tools useful for collecting test samples (such as blood, urine, saliva and stool). Such tools include lancets and absorbent paper or cloth for collecting and stabilizing blood; swabs for collecting and stabilizing saliva; and cups for collecting and stabilizing urine or stool samples. Collection materials such as, papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. These collection materials also may be treated with or contain preservatives, stabilizers or antimicrobial agents to help maintain the integrity of the specimens. Also, the polypeptide can be attached to a solid phase.

Another assay kit for determining the presence of BS106 antigen or anti-BS106 antibody in a test sample comprises a container containing an antibody which specifically binds to a BS106 antigen, wherein the BS106 antigen comprises at least one BS106-encoded epitope. The BS106 antigen has at least about 60% sequence similarity to a sequence of a BS106-encoded antigen selected from the group

consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof. These test kits can further comprise containers with tools useful for collecting test samples (such as blood, urine, saliva and stool). Such tools include lancets and absorbent  
5 paper or cloth for collecting and stabilizing blood; swabs for collecting and stabilizing saliva; cups for collecting and stabilizing urine or stool samples. Collection materials, papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. These collection materials also may be treated with, or contain, preservatives, stabilizers or  
10 antimicrobial agents to help maintain the integrity of the specimens. The antibody can be attached to a solid phase.

A method for producing a polypeptide which contains at least one epitope of BS106 is provided, which method comprises incubating host cells transfected with an expression vector. This vector comprises a polynucleotide sequence encoding a  
15 polypeptide, wherein the polypeptide comprises an amino acid sequence having at least 50% identity to a BS106 amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

A method for detecting BS106 antigen in a test sample suspected of  
20 containing BS106 antigen also is provided. The method comprises contacting the test sample with an antibody or fragment thereof which specifically binds to at least one epitope of a BS106 antigen, for a time and under conditions sufficient for the formation of antibody/antigen complexes; and detecting the presence of such complexes containing the antibody as an indication of the presence of BS106 antigen  
25 in the test sample. The antibody can be attached to a solid phase and be either a monoclonal or polyclonal antibody. Furthermore, the antibody specifically binds to at least one BS106 antigen selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

30 Another method is provided which detects antibodies which specifically bind to BS106 antigen in a test sample suspected of containing these antibodies. The method comprises contacting the test sample with a polypeptide which contains at least one BS106 epitope, wherein the BS106 epitope comprises an amino acid sequence having at least 50% identity with an amino acid sequence encoded by a  
35 BS106 polynucleotide, or a fragment thereof. Contacting is carried out for a time

and under conditions sufficient to allow antigen/antibody complexes to form. The method further entails detecting complexes which contain the polypeptide. The polypeptide can be attached to a solid phase. Further, the polypeptide can be a recombinant protein or a synthetic peptide having at least 50% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16,  
5 SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

The present invention provides a cell transfected with a BS106 nucleic acid sequence that encodes at least one epitope of a BS106 antigen, or fragment thereof.  
10 The nucleic acid sequence is selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.

A method for producing antibodies to BS106 antigen also is provided, which method comprises administering to an individual an isolated immunogenic  
15 polypeptide or fragment thereof, wherein the isolated immunogenic polypeptide comprises at least one BS106 epitope in an amount sufficient to produce an immune response. The isolated, immunogenic polypeptide comprises an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19,  
20 SEQUENCE ID NO 20, and fragments thereof.

Another method for producing antibodies which specifically bind to BS106 antigen is disclosed, which method comprises administering to a mammal a plasmid comprising a nucleic acid sequence which encodes at least one BS106 epitope derived from an amino acid sequence selected from the group consisting of  
25 SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

Also provided is a composition of matter that comprises a BS106 polynucleotide of at least about 10-12 nucleotides having at least 50% identity to a polynucleotide selected from the group consisting of SEQUENCE ID NO 1,  
30 SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof. The BS106 polynucleotide encodes an amino acid sequence having at least one BS106 epitope. Another composition of matter provided by the present invention comprises a polypeptide with at least one BS106 epitope of about 8-10 amino acids. The polypeptide  
35 comprises an amino acid sequence having at least 50% identity to an amino acid

sequence selected from the group consisting of SEQUENCE ID NO 16,  
SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19,  
SEQUENCE ID NO 20, and fragments of SEQUENCE ID NOS 17-20. Also  
provided is a gene, or fragment thereof, coding for a BS106 polypeptide which has  
5 at least 50% identity to SEQUENCE ID NO 16; and a gene, or a fragment thereof,  
comprising DNA having at least 50% identity to SEQUENCE ID NO 4 or  
SEQUENCE ID NO 5.

#### Brief Description of the Drawings

10 FIGURE 1 shows the nucleotide alignment of clones 1662885 (SEQUENCE  
ID NO 1), 893988 (SEQUENCE ID NO 2), 1209814 (SEQUENCE ID NO 3) and  
the consensus sequence (SEQUENCE ID NO 4) derived therefrom. A G/T  
polymorphism at position 54 of the consensus sequence is denoted as "K," and a  
C/T polymorphism at position 312 of the consensus sequence is denoted as "Y."

15 FIGURE 2 shows the contig map depicting the formation of the consensus  
nucleotide sequence (SEQUENCE ID NO 4) from the nucleotide alignment of  
overlapping clones 1662885 (SEQUENCE ID NO 1), 893988 (SEQUENCE ID NO  
2), and 1209814 (SEQUENCE ID NO 3).

FIGURE 3A is a scan of an ethidium bromide-stained agarose gel of RNA  
20 from various tissues and the corresponding northern blot of RNA using BS106  
radiolabeled probe; FIGURE 3B is a scan of another ethidium bromide-stained  
agarose gel of RNA from normal breast and breast cancer tissues and the  
corresponding northern blot of RNA using BS106 radiolabeled probe.

FIGURES 4A and 4B are scans of stained agarose gels of BS106-specific  
25 primed PCR amplification products.

#### Detailed Description of the Invention

The present invention provides a gene or a fragment thereof which codes for  
a BS106 polypeptide having at least about 50% identity to SEQUENCE ID NO 16.  
30 The present invention further encompasses a BS106 gene or a fragment thereof  
comprising DNA which has at least about 50% identity to SEQUENCE ID NO 4 or  
SEQUENCE ID NO 5.

The present invention provides methods for assaying a test sample for  
products of a breast tissue gene designated as BS106, which method comprises  
35 making cDNA from mRNA in the test sample and detecting the cDNA as an  
indication of the presence of breast tissue gene BS106. The method may include an

amplification step, wherein one or more portions of the mRNA from BS106 corresponding to the gene or fragments thereof, is amplified. Methods also are provided for assaying for the translation products of BS106. Test samples which may be assayed by the methods provided herein include tissues, cells, body fluids  
5 and secretions. The present invention also provides reagents such as oligonucleotide primers and polypeptides which are useful in performing these methods.

Portions of the nucleic acid sequences disclosed herein are useful as primers for the reverse transcription of RNA or for the amplification of cDNA; or as probes to determine the presence of certain mRNA sequences in test samples. Also  
10 disclosed are nucleic acid sequences which permit the production of encoded polypeptide sequences which are useful as standards or reagents in diagnostic immunoassays, as targets for pharmaceutical screening assays and/or as components or as target sites for various therapies. Monoclonal and polyclonal antibodies directed against at least one epitope contained within these polypeptide sequences are  
15 useful as delivery agents for therapeutic agents as well as for diagnostic tests and for screening for diseases or conditions associated with BS106, especially breast cancer. Isolation of sequences of other portions of the gene of interest can be accomplished utilizing probes or PCR primers derived from these nucleic acid sequences. This allows additional probes of the mRNA or cDNA of interest to be established, as well  
20 as corresponding encoded polypeptide sequences. These additional molecules are useful in detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining the predisposition to, diseases and conditions of the breast such as breast cancer, characterized by BS106, as disclosed herein.

Techniques for determining amino acid sequence "similarity" are well-known  
25 in the art. In general, "similarity" means the exact amino acid to amino acid comparison of two or more polypeptides at the appropriate place, where amino acids are identical or possess similar chemical and/or physical properties such as charge or hydrophobicity. A so-termed "percent similarity" then can be determined between the compared polypeptide sequences. Techniques for determining nucleic acid and  
30 amino acid sequence identity also are well known in the art and include determining the nucleotide sequence of the mRNA for that gene (usually via a cDNA intermediate) and determining the amino acid sequence encoded thereby, and comparing this to a second amino acid sequence. In general, "identity" refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of two  
35 polynucleotides or polypeptide sequences, respectively. Two or more

polynucleotide sequences can be compared by determining their "percent identity." Two or more amino acid sequences likewise can be compared by determining their "percent identity." The programs available in the Wisconsin Sequence Analysis Package, Version 8 (available from Genetics Computer Group, Madison, WI), for example, the GAP program, are capable of calculating both the identity between two polynucleotides and the identity and similarity between two polypeptide sequences, respectively. Other programs for calculating identity or similarity between sequences are known in the art.

The compositions and methods described herein will enable the identification of certain markers as indicative of a breast tissue disease or condition; the information obtained therefrom will aid in the detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining diseases or conditions associated with BS106, especially breast cancer. Test methods include, for example, probe assays which utilize the sequence(s) provided herein and which also may utilize nucleic acid amplification methods such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), and hybridization. In addition, the nucleotide sequences provided herein contain open reading frames from which an immunogenic epitope may be found. This epitope is believed to be unique to the disease state or condition associated with BS106. It also is thought that the polynucleotides or polypeptides and protein encoded by the BS106 gene are useful as a marker. This marker is either elevated in disease such as breast cancer, altered in disease such as breast cancer, or present as a normal protein but appearing in an inappropriate body compartment. The uniqueness of the epitope may be determined by its immunological reactivity and specificity with antibodies directed against proteins and polypeptides encoded by the BS106 gene, and (ii) its nonreactivity with any other tissue markers. Methods for determining immunological reactivity are well-known and include but are not limited to, for example, radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), hemagglutination (HA), fluorescence polarization immunoassay (FPIA), chemiluminescent immunoassay (CLIA) and others. Several examples of suitable methods are described herein.

Unless otherwise stated, the following terms shall have the following meanings:

A polynucleotide "derived from" or "specific for" a designated sequence refers to a polynucleotide sequence which comprises a contiguous sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides,

more preferably at least about 10-12 nucleotides, and even more preferably at least about 15-20 nucleotides corresponding, i.e., identical or complementary to, a region of the designated nucleotide sequence. The sequence may be complementary or identical to a sequence which is unique to a particular polynucleotide sequence as  
5 determined by techniques known in the art. Comparisons to sequences in databanks, for example, can be used as a method to determine the uniqueness of a designated sequence. Regions from which sequences may be derived, include but are not limited to, regions encoding specific epitopes, as well as non-translated and/or non-transcribed regions.

10 The derived polynucleotide will not necessarily be derived physically from the nucleotide sequence of interest under study, but may be generated in any manner, including but not limited to chemical synthesis, replication, reverse transcription or transcription, which is based on the information provided by the sequence of bases in the region(s) from which the polynucleotide is derived. As such, it may represent  
15 either a sense or an antisense orientation of the original polynucleotide. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with the intended use.

A "fragment" of a specified polynucleotide refers to a polynucleotide sequence which comprises a contiguous sequence of approximately at least about 6  
20 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10-12 nucleotides, and even more preferably at least about 15-20 nucleotides corresponding, i.e., identical or complementary to, a region of the specified nucleotide sequence.

The term "primer" denotes a specific oligonucleotide sequence which is  
25 complementary to a target nucleotide sequence and used to hybridize to the target nucleotide sequence. A primer serves as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase.

The term "probe" denotes a defined nucleic acid segment (or nucleotide  
30 analog segment, e.g., PNA as defined hereinbelow) which can be used to identify a specific polynucleotide present in samples bearing the complementary sequence.

"Encoded by" refers to a nucleic acid sequence which codes for a polypeptide  
sequence, wherein the polypeptide sequence or a portion thereof contains an amino  
acid sequence of at least 3 to 5 amino acids, more preferably at least 8 to 10 amino  
35 acids, and even more preferably at least 15 to 20 amino acids from a polypeptide



encoded by the nucleic acid sequence. Also encompassed are polypeptide sequences which are immunologically identifiable with a polypeptide encoded by the sequence. Thus, a "polypeptide," "protein," or "amino acid" sequence has at least about 50% identity, preferably about 60% identity, more preferably about 75-85% identity, and  
5 most preferably about 90-95% or more identity to a BS106 amino acid sequence. Further, the BS106 "polypeptide," "protein," or "amino acid" sequence may have at least about 60% similarity, preferably at least about 75% similarity, more preferably about 85% similarity, and most preferably about 95% or more similarity to a polypeptide or amino acid sequence of BS106. This amino acid sequence can be  
10 selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

A "recombinant polypeptide," "recombinant protein," or "a polypeptide produced by recombinant techniques," which terms may be used interchangeably  
15 herein, describes a polypeptide which by virtue of its origin or manipulation is not associated with all or a portion of the polypeptide with which it is associated in nature and/or is linked to a polypeptide other than that to which it is linked in nature. A recombinant or encoded polypeptide or protein is not necessarily translated from a designated nucleic acid sequence. It also may be generated in any manner, including  
20 chemical synthesis or expression of a recombinant expression system.

The term "synthetic peptide" as used herein means a polymeric form of amino acids of any length, which may be chemically synthesized by methods well-known to the routineer. These synthetic peptides are useful in various applications.

The term "polynucleotide" as used herein means a polymeric form of  
25 nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, the term includes double- and single-stranded DNA, as well as double- and single-stranded RNA. It also includes modifications, such as methylation or capping and unmodified forms of the polynucleotide. The terms "polynucleotide," "oligomer," "oligonucleotide," and  
30 "oligo" are used interchangeably herein.

"A sequence corresponding to a cDNA" means that the sequence contains a polynucleotide sequence that is identical or complementary to a sequence in the designated DNA. The degree (or "percent") of identity or complementarity to the cDNA will be approximately 50% or greater, preferably at least about 70% or  
35 greater, and more preferably at least about 90% or greater. The sequence that

corresponds to the identified cDNA will be at least about 50 nucleotides in length, preferably at least about 60 nucleotides in length, and more preferably at least about 70 nucleotides in length. The correspondence between the gene or gene fragment of interest and the cDNA can be determined by methods known in the art and include, for example, a direct comparison of the sequenced material with the cDNAs described, or hybridization and digestion with single strand nucleases, followed by size determination of the digested fragments.

“Purified polynucleotide” refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

“Purified polypeptide” or “purified protein” means a polypeptide of interest or fragment thereof which is essentially free of, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, cellular components with which the polypeptide of interest is naturally associated. Methods for purifying polypeptides of interest are known in the art.

The term “isolated” means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, which is separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition, and still be isolated in that the vector or composition is not part of its natural environment.

“Polypeptide” and “protein” are used interchangeably herein and indicate at least one molecular chain of amino acids linked through covalent and/or non-covalent bonds. The terms do not refer to a specific length of the product. Thus peptides, oligopeptides and proteins are included within the definition of polypeptide. The terms include post-translational modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. In addition, protein

fragments, analogs, mutated or variant proteins, fusion proteins and the like are included within the meaning of polypeptide.

A "fragment" of a specified polypeptide refers to an amino acid sequence which comprises at least about 3-5 amino acids, more preferably at least about 8-10 amino acids, and even more preferably at least about 15-20 amino acids derived from the specified polypeptide.

"Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vectors or other transferred DNA, and include the original progeny of the original cell which has been transfected.

As used herein "replicon" means any genetic element, such as a plasmid, a chromosome or a virus, that behaves as an autonomous unit of polynucleotide replication within a cell.

A "vector" is a replicon in which another polynucleotide segment is attached, such as to bring about the replication and/or expression of the attached segment.

The term "control sequence" refers to a polynucleotide sequence which is necessary to effect the expression of a coding sequence to which it is ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, such control sequences generally include a promoter, a ribosomal binding site, and terminators; in eukaryotes, such control sequences generally include promoters, terminators and, in some instances, enhancers. The term "control sequence" thus is intended to include at a minimum all components whose presence is necessary for expression, and also may include additional components whose presence is advantageous, for example, leader sequences.

"Operably linked" refers to a situation wherein the components described are in a relationship permitting them to function in their intended manner. Thus, for example, a control sequence "operably linked" to a coding sequence is ligated in such a manner that expression of the coding sequence is achieved under conditions compatible with the control sequence.

The term "open reading frame" or "ORF" refers to a region of a polynucleotide sequence which encodes a polypeptide. This region may represent a portion of a coding sequence or a total coding sequence.

A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and translated into a polypeptide when placed under the control of

appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5' -terminus and a translation stop codon at the 3' -terminus. A coding sequence can include, but is not limited to, mRNA, cDNA and recombinant polynucleotide sequences.

5           The term "immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptide(s) which also are present in and are unique to the designated polypeptide(s). Immunological identity may be determined by antibody binding and/or competition in binding. These techniques are known to the routineer and also are described herein. The uniqueness of an epitope also can be determined  
10 by computer searches of known data banks, such as GenBank, for the polynucleotide sequence which encodes the epitope and by amino acid sequence comparisons with other known proteins.

As used herein, "epitope" means an antigenic determinant of a polypeptide or protein. Conceivably, an epitope can comprise three amino acids in a spatial  
15 conformation which is unique to the epitope. Generally, an epitope consists of at least five such amino acids and more usually, it consists of at least eight to ten amino acids. Methods of examining spatial conformation are known in the art and include, for example, x-ray crystallography and two-dimensional nuclear magnetic resonance.

20           A "conformational epitope" is an epitope that is comprised of specific juxtaposition of amino acids in an immunologically recognizable structure, such amino acids being present on the same polypeptide in a contiguous or non-contiguous order or present on different polypeptides.

A polypeptide is "immunologically reactive" with an antibody when it binds  
25 to an antibody due to antibody recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly, by the kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an epitope against which the antibody is directed. The methods for determining whether a polypeptide  
30 is immunologically reactive with an antibody are known in the art.

As used herein, the term "immunogenic polypeptide containing an epitope of interest" means naturally occurring polypeptides of interest or fragments thereof, as well as polypeptides prepared by other means, for example, by chemical synthesis or the expression of the polypeptide in a recombinant organism.

The term “transfection” refers to the introduction of an exogenous polynucleotide into a prokaryotic or eucaryotic host cell, irrespective of the method used for the introduction. The term “transfection” refers to both stable and transient introduction of the polynucleotide, and encompasses direct uptake of

5 polynucleotides, transformation, transduction, and f-mating. Once introduced into the host cell, the exogenous polynucleotide may be maintained as a non-integrated replicon, for example, a plasmid, or alternatively, may be integrated into the host genome.

“Treatment” refers to prophylaxis and/or therapy.

10 The term “individual” as used herein refers to vertebrates, particularly members of the mammalian species and includes, but is not limited to, domestic animals, sports animals, primates and humans; more particularly the term refers to humans.

The term “sense strand” or “plus strand” (or “+”) as used herein denotes a

15 nucleic acid that contains the sequence that encodes the polypeptide. The term “antisense strand” or “minus strand” (or “-”) denotes a nucleic acid that contains a sequence that is complementary to that of the “plus” strand.

The term “test sample” refers to a component of an individual's body which is the source of the analyte (such as, antibodies of interest or antigens of interest).

20 These components are well known in the art. A test sample is typically anything suspected of containing a target sequence. Test samples can be prepared using methodologies well known in the art such as by obtaining a specimen from an individual and, if necessary, disrupting any cells contained thereby to release target nucleic acids. These test samples include biological samples which can be tested by

25 the methods of the present invention described herein and include human and animal body fluids such as whole blood, serum, plasma, cerebrospinal fluid, sputum, bronchial washing, bronchial aspirates, urine, lymph fluids and various external secretions of the respiratory, intestinal and genitourinary tracts, tears, saliva, milk, white blood cells, myelomas and the like; biological fluids such as cell culture

30 supernatants; tissue specimens which may be fixed; and cell specimens which may be fixed.

“Purified product” refers to a preparation of the product which has been isolated from the cellular constituents with which the product is normally associated and from other types of cells which may be present in the sample of interest.

“TEST” 16029680

“PNA” denotes a “peptide nucleic acid analog” which may be utilized in a procedure such as an assay described herein to determine the presence of a target. “MA” denotes a “morpholino analog” which may be utilized in a procedure such as an assay described herein to determine the presence of a target. See, for example, 5 U.S. Patent No. 5,378,841, which is incorporated herein by reference. PNAs are neutrally charged moieties which can be directed against RNA targets or DNA. PNA probes used in assays in place of, for example, the DNA probes of the present invention, offer advantages not achievable when DNA probes are used. These advantages include manufacturability, large scale labeling, reproducibility, stability, 10 insensitivity to changes in ionic strength and resistance to enzymatic degradation which is present in methods utilizing DNA or RNA. These PNAs can be labeled with (“attached to”) such signal generating compounds as fluorescein, radionucleotides, chemiluminescent compounds and the like. PNAs or other nucleic acid analogs such as MAs thus can be used in assay methods in place of DNA or 15 RNA. Although assays are described herein utilizing DNA probes, it is within the scope of the routine that PNAs or MAs can be substituted for RNA or DNA with appropriate changes if and as needed in assay reagents.

“Analyte,” as used herein, is the substance to be detected which may be present in the test sample. The analyte can be any substance for which there exists a 20 naturally occurring specific binding member (such as, an antibody), or for which a specific binding member can be prepared. Thus, an analyte is a substance that can bind to one or more specific binding members in an assay. “Analyte” also includes any antigenic substances, haptens, antibodies and combinations thereof. As a member of a specific binding pair, the analyte can be detected by means of naturally 25 occurring specific binding partners (pairs) such as the use of intrinsic factor protein as a member of a specific binding pair for the determination of Vitamin B12, the use of folate-binding protein to determine folic acid, or the use of a lectin as a member of a specific binding pair for the determination of a carbohydrate. The analyte can include a protein, a polypeptide, an amino acid, a nucleotide target and the like.

30 “Diseases of the breast” or “breast disease,” or “condition of the breast,” as used herein, refer to any disease or condition of the breast including, but not limited to, atypical hyperplasia, fibroadenoma, cystic breast disease, and cancer.

“Breast cancer,” as used herein, refers to any malignant disease of the breast including, but not limited to, ductal carcinoma in situ, lobular carcinoma in situ, 35 infiltrating ductal carcinoma, medullary carcinoma, tubular carcinoma, mucinous



be a member of any specific binding pair, including either hapten-anti-hapten systems such as biotin or anti-biotin, avidin or biotin, a carbohydrate or a lectin, a complementary nucleotide sequence, an effector or a receptor molecule, an enzyme cofactor and an enzyme, an enzyme inhibitor or an enzyme and the like. An  
5 immunoreactive specific binding member can be an antibody, an antigen, or an antibody/antigen complex that is capable of binding either to the polypeptide of interest as in a sandwich assay, to the capture reagent as in a competitive assay, or to the ancillary specific binding member as in an indirect assay. When describing probes and probe assays, the term "reporter molecule" may be used. A reporter  
10 molecule comprises a signal generating compound as described hereinabove conjugated to a specific binding member of a specific binding pair, such as carbazole or adamantane.

The various "signal-generating compounds" (labels) contemplated include chromagens, catalysts such as enzymes, luminescent compounds such as fluorescein  
15 and rhodamine, chemiluminescent compounds such as dioxetanes, acridiniums, phenanthridiniums and luminol, radioactive elements and direct visual labels. Examples of enzymes include alkaline phosphatase, horseradish peroxidase, beta-galactosidase and the like. The selection of a particular label is not critical, but it must be capable of producing a signal either by itself or in conjunction with one or  
20 more additional substances.

"Solid phases" ("solid supports") are known to those in the art and include the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic or non-magnetic beads, nitrocellulose strips, membranes, microparticles such as latex  
25 particles, sheep (or other animal) red blood cells and Duracytes® (red blood cells "fixed" by pyruvic aldehyde and formaldehyde, available from Abbott Laboratories, Abbott Park, IL) and others. The "solid phase" is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon  
30 chips, sheep (or other suitable animal's) red blood cells and Duracytes® are all suitable examples. Suitable methods for immobilizing peptides on solid phases include ionic, hydrophobic, covalent interactions and the like. A "solid phase," as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid phase can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an  
35 additional receptor which has the ability to attract and immobilize the capture reagent.



The additional receptor can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid phase and which  
5 has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid phase material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead,  
10 microparticle, chip, sheep (or other suitable animal's) red blood cells, Duracytes® and other configurations known to those of ordinary skill in the art.

It is contemplated and within the scope of the present invention that the solid phase also can comprise any suitable porous material with sufficient porosity to allow access by detection antibodies and a suitable surface affinity to bind antigens.  
15 Microporous structures generally are preferred, but materials with a gel structure in the hydrated state may be used as well. Such useful solid supports include, but are not limited to, nitrocellulose and nylon. It is contemplated that such porous solid supports described herein preferably are in the form of sheets of thickness from about 0.01 to 0.5 mm, preferably about 0.1 mm. The pore size may vary within  
20 wide limits and preferably is from about 0.025 to 15 microns, especially from about 0.15 to 15 microns. The surface of such supports may be activated by chemical processes which cause covalent linkage of the antigen or antibody to the support. The irreversible binding of the antigen or antibody is obtained, however, in general, by adsorption on the porous material by poorly understood hydrophobic forces.  
25 Other suitable solid supports are known in the art.

#### Reagents

The present invention provides reagents such as polynucleotide sequences derived from a breast tissue of interest and designated as BS106, polypeptides encoded thereby and antibodies specific for these polypeptides. The present  
30 invention also provides reagents such as oligonucleotide fragments derived from the disclosed polynucleotides and nucleic acid sequences complementary to these polynucleotides. The polynucleotides, polypeptides, or antibodies of the present invention may be used to provide information leading to the detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating of, or determining the  
35 predisposition to, diseases and conditions of the breast such as cancer. The

sequences disclosed herein represent unique polynucleotides which can be used in assays or for producing a specific profile of gene transcription activity. Such assays are disclosed in European Patent Number 0373203B1 and International Publication No. WO 95/11995, which are hereby incorporated by reference.

5           Selected BS106-derived polynucleotides can be used in the methods described herein for the detection of normal or altered gene expression. Such methods may employ BS106 polynucleotides or oligonucleotides, fragments or derivatives thereof, or nucleic acid sequences complementary thereto.

10           The polynucleotides disclosed herein, their complementary sequences, or fragments of either, can be used in assays to detect, amplify or quantify genes, nucleic acids, cDNAs or mRNAs relating to breast tissue disease and conditions associated therewith. They also can be used to identify an entire or partial coding region of a BS106 polypeptide. They further can be provided in individual containers in the form of a kit for assays, or provided as individual compositions. If  
15           provided in a kit for assays, other suitable reagents such as buffers, conjugates and the like may be included.

20           The polynucleotide may be in the form of RNA or DNA. Polynucleotides in the form of DNA, cDNA, genomic DNA, nucleic acid analogs and synthetic DNA are within the scope of the present invention. The DNA may be double-stranded or single-stranded, and if single stranded, may be the coding (sense) strand or non-coding (anti-sense) strand. The coding sequence which encodes the polypeptide may be identical to the coding sequence provided herein or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same polypeptide as the DNA provided herein.

25           This polynucleotide may include only the coding sequence for the polypeptide, or the coding sequence for the polypeptide and an additional coding sequence such as a leader or secretory sequence or a proprotein sequence, or the coding sequence for the polypeptide (and optionally an additional coding sequence) and a non-coding sequence, such as a non-coding sequence 5' and/or 3' of the  
30           coding sequence for the polypeptide.

35           In addition, the invention includes variant polynucleotides containing modifications such as polynucleotide deletions, substitutions or additions; and any polypeptide modification resulting from the variant polynucleotide sequence. A polynucleotide of the present invention also may have a coding sequence which is a naturally occurring allelic variant of the coding sequence provided herein.

In addition, the coding sequence for the polypeptide may be fused in the same reading frame to a polynucleotide sequence which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the polypeptide. The polynucleotides may also encode for a proprotein which is the protein plus additional 5' amino acid residues. A protein having a prosequence is a proprotein and may, in some cases, be an inactive form of the protein. Once the prosequence is cleaved an active protein remains. Thus, the polynucleotide of the present invention may encode for a protein, or for a protein having a prosequence, or for a protein having both a prosequence (leader sequence) and a prosequence.

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g. a COS-7 cell line, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein. See, for example, I. Wilson et al., Cell 37:767 (1984).

It is contemplated that polynucleotides will be considered to hybridize to the sequences provided herein if there is at least 50%, preferably at least 70%, and more preferably at least 90% identity between the polynucleotide and the sequence.

The present invention also provides an antibody produced by using a purified BS106 polypeptide of which at least a portion of the polypeptide is encoded by a BS106 polynucleotide selected from the polynucleotides provided herein. These antibodies may be used in the methods provided herein for the detection of BS106 antigen in test samples. The presence of BS106 antigen in the test samples is indicative of the presence of a breast disease or condition. The antibody also may be used for therapeutic purposes, for example, in neutralizing the activity of BS106 polypeptide in conditions associated with altered or abnormal expression.

The present invention further relates to a BS106 polypeptide which has the deduced amino acid sequence as provided herein, as well as fragments, analogs and derivatives of such polypeptide. The polypeptide of the present invention may be a recombinant polypeptide, a natural purified polypeptide or a synthetic polypeptide.

46TCT 1609680

The fragment, derivative or analog of the BS106 polypeptide may be one in which one or more of the amino acid residues is substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code; 5 or it may be one in which one or more of the amino acid residues includes a substituent group; or it may be one in which the polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol); or it may be one in which the additional amino acids are fused to the polypeptide, such as a leader or secretory sequence or a sequence 10 which is employed for purification of the polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are within the scope of the present invention. The polypeptides and polynucleotides of the present invention are provided preferably in an isolated form and preferably purified.

Thus, a polypeptide of the present invention may have an amino acid 15 sequence that is identical to that of the naturally occurring polypeptide or that is different by minor variations due to one or more amino acid substitutions. The variation may be a "conservative change" typically in the range of about 1 to 5 amino acids, wherein the substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine or threonine with serine. In 20 contrast, variations may include nonconservative changes, e.g., replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without changing biological or immunological activity may be found using computer programs well 25 known in the art, for example, DNASTAR software (DNASTAR Inc., Madison WI).

Probes constructed according to the polynucleotide sequences of the present invention can be used in various assay methods to provide various types of analysis. For example, such probes can be used in fluorescent in situ hybridization (FISH) 30 technology to perform chromosomal analysis, and used to identify cancer-specific structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR-generated and/or allele specific oligonucleotides probes, allele specific amplification or by direct sequencing. Probes also can be labeled with radioisotopes, directly- or indirectly- 35 detectable haptens, or fluorescent molecules, and utilized for in situ hybridization

studies to evaluate the mRNA expression of the gene comprising the polynucleotide in tissue specimens or cells.

This invention also provides teachings as to the production of the polynucleotides and polypeptides provided herein.

5        Probe Assays

The sequences provided herein may be used to produce probes which can be used in assays for the detection of nucleic acids in test samples. The probes may be designed from conserved nucleotide regions of the polynucleotides of interest or from non-conserved nucleotide regions of the polynucleotide of interest. The design  
10 of such probes for optimization in assays is within the skill of the routineer. Generally, nucleic acid probes are developed from non-conserved or unique regions when maximum specificity is desired, and nucleic acid probes are developed from conserved regions when assaying for nucleotide regions that are closely related to, for example, different members of a multi-gene family or in related species like  
15 mouse and man.

The polymerase chain reaction (PCR) is a technique for amplifying a desired nucleic acid sequence (target) contained in a nucleic acid or mixture thereof. In PCR, a pair of primers are employed in excess to hybridize to the complementary strands of the target nucleic acid. The primers are each extended by a polymerase  
20 using the target nucleic acid as a template. The extension products become target sequences themselves, following dissociation from the original target strand. New primers then are hybridized and extended by a polymerase, and the cycle is repeated to geometrically increase the number of target sequence molecules. PCR is disclosed in U.S. Patents 4,683,195 and 4,683,202, which are incorporated herein by  
25 reference.

The Ligase Chain Reaction (LCR) is an alternate method for nucleic acid amplification. In LCR, probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target  
30 strand, and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3' hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can  
35 hybridize to a portion of the second probe in a similar abutting fashion. Of course, if

the target is initially double stranded, the secondary probes also will hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes which can be ligated to form a complementary, secondary ligated product. It is  
5 important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization and ligation, amplification of the target sequence is achieved. This technique is described more completely in EP-A- 320 308 to K. Backman published June 16, 1989 and EP-A- 439 182 to K. Backman et al, published July 31, 1991, both of which are  
10 incorporated herein by reference.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Patent No. 5,322,770, which is incorporated herein by reference; or reverse transcribe mRNA  
15 into cDNA followed by asymmetric gap ligase chain reaction (RT-AGLCR) as described by R.L. Marshall et al., PCR Methods and Applications 4: 80-84 (1994), which also is incorporated herein by reference.

Other known amplification methods which can be utilized herein include but are not limited to the so-called "NASBA" or "3SR" technique described by J.C. Guatelli, et al., PNAS USA 87:1874-1878 (1990) and also described by J. Compton, Nature 350 (No. 6313):91-92 (1991); Q-beta amplification as described in published European Patent Application (EPA) No. 4544610; strand displacement amplification (as described in G.T. Walker et al., Clin. Chem. 42:9-13 (1996)) and  
20 European Patent Application No. 684315; and target mediated amplification, as described in International Publication No. WO 93/22461.

Detection of BS106 may be accomplished using any suitable detection method, including those detection methods which are currently well known in the art, as well as detection strategies which may evolve later. Examples of the foregoing presently known detection methods are hereby incorporated herein by  
30 reference. See, for example, Caskey et al., U.S. Patent No. 5,582,989, Gelfand et al., U.S. Patent No. 5,210,015. Examples of such detection methods include target amplification methods as well as signal amplification technologies. An example of presently known detection methods would include the nucleic acid amplification technologies referred to as PCR, LCR, NASBA, SDA, RCR and TMA. See, for  
35 example, Caskey et al., U.S. Patent No. 5,582,989, Gelfand et al., U.S. Patent No.

0896094-037

5,210,015. All of the foregoing are hereby incorporated by reference. Detection may also be accomplished using signal amplification such as that disclosed in Snitman et al., U.S. Patent No. 5,273,882. While the amplification of target or signal is preferred at present, it is contemplated and within the scope of the present invention that ultrasensitive detection methods which do not require amplification can be utilized herein.

Detection, both amplified and non-amplified, may be (combined) carried out using a variety of heterogeneous and homogeneous detection formats. Examples of heterogeneous detection formats are disclosed in Snitman et al., U.S. Patent No. 5,273,882, Albarella et al in EP-84114441.9, Urdea et al., U.S. Patent No. 5,124,246, Ullman et al. U.S. Patent No. 5,185,243 and Kourilsky et al., U.S. Patent No. 4,581,333. All of the foregoing are hereby incorporated by reference. Examples of homogeneous detection formats are disclosed in, Caskey et al., U.S. Patent No. 5,582,989, Gelfand et al., U.S. Patent No. 5,210,015, which are incorporated herein by reference. Also contemplated and within the scope of the present invention is the use of multiple probes in the hybridization assay, which use improves sensitivity and amplification of the BS106 signal. See, for example, Caskey et al., U.S. Patent No. 5,582,989, Gelfand et al., U.S. Patent No. 5,210,015, which are incorporated herein by reference.

In one embodiment, the present invention generally comprises the steps of contacting a test sample suspected of containing a target polynucleotide sequence with amplification reaction reagents comprising an amplification primer, and a detection probe that can hybridize with an internal region of the amplicon sequences. Probes and primers employed according to the method provided herein are labeled with capture and detection labels, wherein probes are labeled with one type of label and primers are labeled with another type of label. Additionally, the primers and probes are selected such that the probe sequence has a lower melt temperature than the primer sequences. The amplification reagents, detection reagents and test sample are placed under amplification conditions whereby, in the presence of target sequence, copies of the target sequence (an amplicon) are produced. In the usual case, the amplicon is double stranded because primers are provided to amplify a target sequence and its complementary strand. The double stranded amplicon then is thermally denatured to produce single stranded amplicon members. Upon formation of the single stranded amplicon members, the mixture is cooled to allow the formation of complexes between the probes and single stranded amplicon members.

As the single stranded amplicon sequences and probe sequences are cooled, the probe sequences preferentially bind the single stranded amplicon members. This finding is counterintuitive given that the probe sequences generally are selected to be shorter than the primer sequences and therefore have a lower melt temperature than the primers. Accordingly, the melt temperature of the amplicon produced by the primers should also have a higher melt temperature than the probes. Thus, as the mixture cools, the re-formation of the double stranded amplicon would be expected. As previously stated, however, this is not the case. The probes are found to preferentially bind the single stranded amplicon members. Moreover, this preference of probe/single stranded amplicon binding exists even when the primer sequences are added in excess of the probes.

After the probe/single stranded amplicon member hybrids are formed, they are detected. Standard heterogeneous assay formats are suitable for detecting the hybrids using the detection labels and capture labels present on the primers and probes. The hybrids can be bound to a solid phase reagent by virtue of the capture label and detected by virtue of the detection label. In cases where the detection label is directly detectable, the presence of the hybrids on the solid phase can be detected by causing the label to produce a detectable signal, if necessary, and detecting the signal. In cases where the label is not directly detectable, the captured hybrids can be contacted with a conjugate, which generally comprises a binding member attached to a directly detectable label. The conjugate becomes bound to the complexes and the conjugates presence on the complexes can be detected with the directly detectable label. Thus, the presence of the hybrids on the solid phase reagent can be determined. Those skilled in the art will recognize that wash steps may be employed to wash away unhybridized amplicon or probe as well as unbound conjugate.

Although the target sequence is described as single stranded, it also is contemplated to include the case where the target sequence is actually double stranded but is merely separated from its complement prior to hybridization with the amplification primer sequences. In the case where PCR is employed in this method, the ends of the target sequences are usually known. In cases where LCR or a modification thereof is employed in the preferred method, the entire target sequence is usually known. Typically, the target sequence is a nucleic acid sequence such as, for example, RNA or DNA.

The method provided herein can be used in well-known amplification reactions that include thermal cycle reaction mixtures, particularly in PCR and gap



LCR (GLCR). Amplification reactions typically employ primers to repeatedly generate copies of a target nucleic acid sequence, which target sequence is usually a small region of a much larger nucleic acid sequence. Primers are themselves nucleic acid sequences that are complementary to regions of a target sequence. Under  
5 amplification conditions, these primers hybridize or bind to the complementary regions of the target sequence. Copies of the target sequence typically are generated by the process of primer extension and/or ligation which utilizes enzymes with polymerase or ligase activity, separately or in combination, to add nucleotides to the hybridized primers and/or ligate adjacent probe pairs. The nucleotides that are added  
10 to the primers or probes, as monomers or preformed oligomers, are also complementary to the target sequence. Once the primers or probes have been sufficiently extended and/or ligated, they are separated from the target sequence, for example, by heating the reaction mixture to a "melt temperature" which is one in which complementary nucleic acid strands dissociate. Thus, a sequence  
15 complementary to the target sequence is formed.

A new amplification cycle then can take place to further amplify the number of target sequences by separating any double stranded sequences, allowing primers or probes to hybridize to their respective targets, extending and/or ligating the hybridized primers or probes and re-separating. The complementary sequences that  
20 are generated by amplification cycles can serve as templates for primer extension or filling the gap of two probes to further amplify the number of target sequences. Typically, a reaction mixture is cycled between 20 and 100 times, more typically, a reaction mixture is cycled between 25 and 50 times. The numbers of cycles can be determined by the routineer. In this manner, multiple copies of the target sequence  
25 and its complementary sequence are produced. Thus, primers initiate amplification of the target sequence when it is present under amplification conditions.

Generally, two primers which are complementary to a portion of a target strand and its complement are employed in PCR. For LCR, four probes, two of which are complementary to a target sequence and two of which are similarly  
30 complementary to the target's complement, are generally employed. In addition to the primer sets and enzymes previously mentioned, a nucleic acid amplification reaction mixture may also comprise other reagents which are well known and include but are not limited to: enzyme cofactors such as manganese; magnesium; salts; nicotinamide adenine dinucleotide (NAD); and deoxynucleotide triphosphates

2025-10-14 16:09:00

(dNTPs) such, as for example, deoxyadenine triphosphate, deoxyguanine triphosphate, deoxycytosine triphosphate and deoxythymine triphosphate.

While the amplification primers initiate amplification of the target sequence, the detection (or hybridization) probe is not involved in amplification. Detection  
5 probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example, peptide nucleic acids which are disclosed in International Publication No. WO 92/20702; morpholino analogs which are described in U.S. Patents Nos 5,185,444, 5,034,506 and 5,142,047; and the like. Depending upon the type of label carried by the probe, the probe is employed to capture or detect the  
10 amplicon generated by the amplification reaction. The probe is not involved in amplification of the target sequence and therefore may have to be rendered "non-extendible" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendible and nucleic acid probes can be rendered non-extendible by modifying the 3' end of the probe such that the hydroxyl  
15 group is no longer capable of participating in elongation. For example, the 3' end of the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified. U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 and incorporated herein by reference describes  
20 modifications which can be used to render a probe non-extendible.

The ratio of primers to probes is not important. Thus, either the probes or primers can be added to the reaction mixture in excess whereby the concentration of one would be greater than the concentration of the other. Alternatively, primers and probes can be employed in equivalent concentrations. Preferably, however, the  
25 primers are added to the reaction mixture in excess of the probes. Thus, primer to probe ratios of, for example, 5:1 and 20:1 are preferred.

While the length of the primers and probes can vary, the probe sequences are selected such that they have a lower melt temperature than the primer sequences. Hence, the primer sequences are generally longer than the probe sequences.  
30 Typically, the primer sequences are in the range of between 20 and 50 nucleotides long, more typically in the range of between 20 and 30 nucleotides long. The typical probe is in the range of between 10 and 25 nucleotides long.

Various methods for synthesizing primers and probes are well known in the art. Similarly, methods for attaching labels to primers or probes are also well known  
35 in the art. For example, it is a matter of routine to synthesize desired nucleic acid

primers or probes using conventional nucleotide phosphoramidite chemistry and instruments available from Applied Biosystems, Inc., (Foster City, CA), DuPont (Wilmington, DE), or Milligen (Bedford MA). Many methods have been described for labeling oligonucleotides such as the primers or probes of the present invention.

5 Enzo Biochemical (New York, NY) and Clontech (Palo Alto, CA) both have described and commercialized probe labeling techniques. For example, a primary amine can be attached to a 3' oligo terminus using 3'-Amine-ON CPG™ (Clontech, Palo Alto, CA). Similarly, a primary amine can be attached to a 5' oligo terminus using Aminomodifier II® (Clontech). The amines can be reacted to various haptens

10 using conventional activation and linking chemistries. In addition, copending applications U.S. Serial Nos. 625,566, filed December 11, 1990 and 630,908, filed December 20, 1990, which are each incorporated herein by reference, teach methods for labeling probes at their 5' and 3' termini, respectively. International Publication Nos WO 92/10505, published 25 June 1992, and WO 92/11388, published 9 July

15 1992, teach methods for labeling probes at their 5' and 3' ends, respectively. According to one known method for labeling an oligonucleotide, a label-phosphoramidite reagent is prepared and used to add the label to the oligonucleotide during its synthesis. See, for example, N.T. Thuong et al., Tet. Letters 29(46):5905-5908 (1988); or J.S. Cohen et al., published U.S. Patent Application

20 07/246,688 (NTIS ORDER No. PAT-APPL-7-246,688) (1989). Preferably, probes are labeled at their 3' and 5' ends.

A capture label is attached to the primers or probes and can be a specific binding member which forms a binding pair with the solid phase reagent's specific binding member. It will be understood that the primer or probe itself may serve as

25 the capture label. For example, in the case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of the primer or probe to thereby immobilize the primer or probe to the solid phase. In cases where the probe itself serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence

30 or "tail" that is not complementary to the single stranded amplicon members. In the case where the primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase because the probe is selected such that it is not fully complementary to the primer sequence.

Generally, probe/single stranded amplicon member complexes can be

35 detected using techniques commonly employed to perform heterogeneous

immunoassays. Preferably, in this embodiment, detection is performed according to the protocols used by the commercially available Abbott LCx<sup>®</sup> instrumentation (Abbott Laboratories, Abbott Park, IL).

The primers and probes disclosed herein are useful in typical PCR assays, wherein the test sample is contacted with a pair of primers, amplification is performed, the hybridization probe is added, and detection is performed.

Another method provided by the present invention comprises contacting a test sample with a plurality of polynucleotides, wherein at least one polynucleotide is a BS106 molecule as described herein, hybridizing the test sample with the plurality of polynucleotides and detecting hybridization complexes. Hybridization complexes are identified and quantitated to compile a profile which is indicative of breast tissue disease, such as breast cancer. Expressed RNA sequences may further be detected by reverse transcription and amplification of the DNA product by procedures well-known in the art, including polymerase chain reaction (PCR).

#### Drug Screening and Gene Therapy.

The present invention also encompasses the use of gene therapy methods for the introduction of anti-sense BS106 derived molecules, such as polynucleotides or oligonucleotides of the present invention, into patients with conditions associated with abnormal expression of polynucleotides related to a breast tissue disease or condition especially breast cancer. These molecules, including antisense RNA and DNA fragments and ribozymes, are designed to inhibit the translation of BS106-mRNA, and may be used therapeutically in the treatment of conditions associated with altered or abnormal expression of a BS106 polynucleotide.

Alternatively, the oligonucleotides described above can be delivered to cells by procedures known in the art such that the anti-sense RNA or DNA may be expressed in vivo to inhibit production of a BS106 polypeptide in the manner described above. Antisense constructs to a BS106 polynucleotide, therefore, reverse the action of BS106 transcripts and may be used for treating breast tissue disease conditions, such as breast cancer. These antisense constructs may also be used to treat tumor metastases.

The present invention also provides a method of screening a plurality of compounds for specific binding to BS106 polypeptide(s), or any fragment thereof, to identify at least one compound which specifically binds the BS106 polypeptide. Such a method comprises the steps of providing at least one compound; combining the BS106 polypeptide with each compound under suitable conditions for a time





employing computer modeling techniques in place of, or in addition to, x-ray crystallography.

Antibodies specific to a BS106 polypeptide (e.g., anti-BS106 antibodies) further may be used to inhibit the biological action of the polypeptide by binding to the polypeptide. In this manner, the antibodies may be used in therapy, for example, to treat breast tissue diseases including breast cancer and its metastases.

Further, such antibodies can detect the presence or absence of the BS106 polypeptide in a test sample and, therefore, are useful as diagnostic markers for the diagnosis of a breast tissue disease or condition especially breast cancer. Such antibodies may also function as a diagnostic marker for breast tissue disease conditions such as breast cancer. The present invention also is directed to antagonists and inhibitors of the polypeptides of the present invention. The antagonists and inhibitors are those which inhibit or eliminate the function of the polypeptide. Thus, for example, an antagonist may bind to a polypeptide of the present invention and inhibit or eliminate its function. The antagonist, for example, could be an antibody against the polypeptide which eliminates the activity of the BS106 polypeptide by binding the BS106 polypeptide, or in some cases the antagonist may be an oligonucleotide. Examples of small molecule inhibitors include, but are not limited to, small peptides or peptide-like molecules.

The antagonists and inhibitors may be employed as a composition with a pharmaceutically acceptable carrier, including, but not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. Administration of BS106 polypeptide inhibitors is preferably systemic. The present invention also provides an antibody which inhibits the action of such a polypeptide.

Antisense technology can be used to reduce gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes for the polypeptide of the present invention, is used to design an antisense RNA oligonucleotide of from 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription, thereby preventing transcription and the production of the BS106 polypeptide. For triple helix, see, for example, Lee et al, Nuc. Acids Res. 6:3073 (1979); Cooney et al, Science 241:456 (1988); and Dervan et al, Science 251:1360 (1991) The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of a mRNA molecule into the BS106

polypeptide. For antisense, see, for example, Okano, J. Neurochem. 56:560 (1991); and "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression," CRC Press, Boca Raton, Fla. (1988). Antisense oligonucleotides act with greater efficacy when modified to contain artificial internucleotide linkages which render the molecule resistant to nucleolytic cleavage. Such artificial internucleotide linkages include, but are not limited to, methylphosphonate, phosphorothiolate and phosphoroamidate internucleotide linkages.

Recombinant Technology.

The present invention provides host cells and expression vectors comprising BS106 polynucleotides of the present invention and methods for the production of the polypeptide(s) they encode. Such methods comprise culturing the host cells under conditions suitable for the expression of the BS106 polynucleotide and recovering the BS106 polypeptide from the cell culture.

The present invention also provides vectors which include BS106 polynucleotides of the present invention, host cells which are genetically engineered with vectors of the present invention and the production of polypeptides of the present invention by recombinant techniques.

Host cells are genetically engineered (transfected, transduced or transformed) with the vectors of this invention which may be cloning vectors or expression vectors. The vector may be in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transfected cells, or amplifying BS106 gene(s). The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing a polypeptide by recombinant techniques. Thus, the polynucleotide sequence may be included in any one of a variety of expression vehicles, in particular vectors or plasmids for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus and pseudorabies. However, any other plasmid or vector may be used so long as it is replicable and viable in the host.



5 The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into appropriate restriction endonuclease sites by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art. The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. Representative examples of such promoters include, but are not limited to, the LTR or the SV40 promoter, the E. coli lac or trp, the phage lambda P sub L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. In addition, the expression vectors preferably contain a gene to provide a phenotypic trait for selection of transfected host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

10 The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transfect an appropriate host to permit the host to express the protein. As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Salmonella typhimurium, Streptomyces sp.; fungal cells, such as yeast; insect cells such as Drosophila and Sf9; animal cells such as CHO, COS or Bowes melanoma; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings provided herein.

15 More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available. The following vectors are provided by way of example. Bacterial: pINCY (Incyte Pharmaceuticals Inc., Palo Alto, CA), pSPORT1 (Life Technologies, Gaithersburg, MD), pQE70, pQE60, pQE-9 (Qiagen) pBs, phagescript, psiX174, pBluescript SK, pBsKS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A, pKK223-3, pKK233-3, pDR540, pRIT5

(Pharmacia); Eukaryotic: pWLneo, pSV2cat, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as it is replicable and viable in the host.

Plasmid pINCY is generally identical to the plasmid pSPORT1 (available  
5 from Life Technologies, Gaithersburg, MD) with the exception that it has two  
modifications in the polylinker (multiple cloning site). These modifications are (1) it  
lacks a HindIII restriction site and (2) its EcoRI restriction site lies at a different  
location. pINCY is created from pSPORT1 by cleaving pSPORT1 with both  
HindIII and EcoRI and replacing the excised fragment of the polylinker with  
10 synthetic DNA fragments (SEQUENCE ID NO 6 and SEQUENCE ID NO 7). This  
replacement may be made in any manner known to those of ordinary skill in the art.  
For example, the two nucleotide sequences, SEQUENCE ID NO 6 and SEQUENCE  
ID NO 7, may be generated synthetically with 5' terminal phosphates, mixed  
together, and then ligated under standard conditions for performing staggered end  
15 ligations into the pSPORT1 plasmid cut with HindIII and EcoRI. Suitable host cells  
(such as *E. coli* DH5 $\alpha$  cells) then are transfected with the ligated DNA and  
recombinant clones are selected for ampicillin resistance. Plasmid DNA then is  
prepared from individual clones and subjected to restriction enzyme analysis or DNA  
sequencing in order to confirm the presence of insert sequences in the proper  
20 orientation. Other cloning strategies known to the ordinary artisan also may be  
employed.

Promoter regions can be selected from any desired gene using CAT  
(chloramphenicol transferase) vectors or other vectors with selectable markers. Two  
appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters  
25 include lacI, lacZ, T3, SP6, T7, gpt, lambda P sub R, P sub L and trp. Eukaryotic  
promoters include cytomegalovirus (CMV) immediate early, herpes simplex virus  
(HSV) thymidine kinase, early and late SV40, LTRs from retroviruses and mouse  
metallothionein-I. Selection of the appropriate vector and promoter is well within the  
level of ordinary skill in the art.

30 In a further embodiment, the present invention provides host cells containing  
the above-described construct. The host cell can be a higher eukaryotic cell, such as  
a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can  
be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the  
host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated  
35 transfection, or electroporation (L. Davis et al., "Basic Methods in Molecular

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicaid   | 58.9%       |
| Private  | 41.1%       |
| Uninsured  | 0.0%        |
| Smoking status   |             |
| Current smoker   | 28.5%       |
| Former smoker  | 35.2%       |
| Never smoker   | 36.3%       |
| Alcohol consumption  |             |
| Regular drinker  | 15.8%       |
| Occasional drinker   | 22.4%       |
| Non-drinker  | 61.8%       |
| Exercise frequency   |             |
| Regular  | 18.9%       |
| Occasional   | 25.6%       |
| Never  | 55.5%       |

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicaid   | 58.9%       |
| Private  | 41.1%       |
| Uninsured  | 0.0%        |
| Smoking status   |             |
| Current smoker   | 28.5%       |
| Former smoker  | 35.2%       |
| Never smoker   | 36.3%       |
| Alcohol consumption  |             |
| Regular drinker  | 15.8%       |
| Occasional drinker   | 22.4%       |
| Non-drinker  | 61.8%       |
| Exercise frequency   |             |
| Regular  | 18.9%       |
| Occasional   | 25.6%       |
| Never  | 55.5%       |

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicaid   | 58.9%       |
| Private  | 41.1%       |
| Uninsured  | 0.0%        |
| Smoking status   |             |
| Current smoker   | 28.5%       |
| Former smoker  | 35.2%       |
| Never smoker   | 36.3%       |
| Alcohol consumption  |             |
| Regular drinker  | 15.8%       |
| Occasional drinker   | 22.4%       |
| Non-drinker  | 61.8%       |
| Exercise frequency   |             |
| Regular  | 18.9%       |
| Occasional   | 25.6%       |
| Never  | 55.5%       |

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicaid   | 58.9%       |
| Private  | 41.1%       |
| Uninsured  | 0.0%        |
| Smoking status   |             |
| Current smoker   | 28.5%       |
| Former smoker  | 35.2%       |
| Never smoker   | 36.3%       |
| Alcohol consumption  |             |
| Regular drinker  | 15.8%       |
| Occasional drinker   | 22.4%       |
| Non-drinker  | 61.8%       |
| Exercise frequency   |             |
| Regular  | 18.9%       |
| Occasional   | 25.6%       |
| Never  | 55.5%       |

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicaid   | 58.9%       |
| Private  | 41.1%       |
| Uninsured  | 0.0%        |
| Smoking status   |             |
| Current smoker   | 28.5%       |
| Former smoker  | 35.2%       |
| Never smoker   | 36.3%       |
| Alcohol consumption  |             |
| Regular drinker  | 15.8%       |
| Occasional drinker   | 22.4%       |
| Non-drinker  | 61.8%       |
| Exercise frequency   |             |
| Regular  | 18.9%       |
| Occasional   | 25.6%       |
| Never  | 55.5%       |

| Table 1. Demographic characteristics of the study population |             |
|--|-------------|
| Age (years)  | Mean (SD)   |
| Male   | 55.2 (10.5) |
| Female   | 56.8 (11.2) |
| Marital status   |             |
| Married  | 78.5%       |
| Single   | 12.3%       |
| Divorced   | 8.2%        |
| Widowed  | 1.0%        |
| Education level  |             |
| High school or less  | 65.4%       |
| College  | 34.6%       |
| Income (US\$)  |             |
| <10,000  | 45.2%       |
| 10,000-20,000  | 32.1%       |
| >20,000  | 22.7%       |
| Health insurance   |             |
| Medicare   | 89.5%       |
| Private  | 10.5%       |
| None   | 0.0%        |
| Comorbidities  |             |
| Hypertension   | 58.3%       |
| Diabetes   | 22.1%       |
| Cholesterol  | 35.7%       |
| Smoking status   |             |
| Current  | 15.2%       |
| Former   | 42.8%       |
| Never  | 42.0%       |
| Alcohol consumption  |             |
| Regular  | 18.5%       |
| Occasional   | 32.1%       |
| None   | 49.4%       |

initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transfection  
5 include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces and Staphylococcus, although, others may also be employed as a routine matter of choice.

Useful expression vectors for bacterial use comprise a selectable marker and bacterial origin of replication derived from plasmids comprising genetic elements of  
10 the well-known cloning vector pBR322 (ATCC 37017). Other vectors include but are not limited to PKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transfection of a suitable host and growth of the host to an  
15 appropriate cell density, the selected promoter is derepressed by appropriate means (e.g., temperature shift or chemical induction), and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Microbial cells employed in expression of proteins can be disrupted by  
20 any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents; such methods are well-known to the ordinary artisan.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-  
25 7 lines of monkey kidney fibroblasts described by Gluzman, Cell 23:175 (1981), and other cell lines capable of expressing a compatible vector, such as the C127, HEK-293, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer and also any necessary ribosome binding sites, polyadenylation sites, splice donor and acceptor  
30 sites, transcriptional termination sequences and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Representative, useful vectors include pRc/CMV and pcDNA3 (available from Invitrogen, San Diego, CA).

BS106 polypeptides are recovered and purified from recombinant cell cultures by known methods including affinity chromatography, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, hydroxyapatite chromatography or lectin chromatography. It is preferred to have low concentrations (approximately 0.1-5 mM) of calcium ion present during purification (Price, et al., J. Biol. Chem. 244:917 (1969)). Protein refolding steps can be used, as necessary, in completing configuration of the polypeptide. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

Thus, polypeptides of the present invention may be naturally purified products expressed from a high expressing cell line, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated with mammalian or other eukaryotic carbohydrates or may be non-glycosylated. The polypeptides of the invention may also include an initial methionine amino acid residue.

The starting plasmids can be constructed from available plasmids in accord with published, known procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

The following is the general procedure for the isolation and analysis of cDNA clones. In a particular embodiment disclosed herein, mRNA was isolated from breast tissue and used to generate the cDNA library. Breast tissue was obtained from patients by surgical resection and was classified as tumor or non-tumor tissue by a pathologist.

The cDNA inserts from random isolates of the breast tissue libraries were sequenced in part, analyzed in detail as set forth in the Examples and are disclosed in the Sequence Listing as SEQUENCE ID NO 1, SEQUENCE ID NO 2, and SEQUENCE ID NO 3. The consensus sequence of these inserts is presented as SEQUENCE ID NO 4. These polynucleotides may contain an entire open reading frame with or without associated regulatory sequences for a particular gene, or they may encode only a portion of the gene of interest. This is attributed to the fact that many genes are several hundred and sometimes several thousand bases in length

and, with current technology, cannot be cloned in their entirety because of vector limitations, incomplete reverse transcription of the first strand, or incomplete replication of the second strand. Contiguous, secondary clones containing additional nucleotide sequences may be obtained using a variety of methods known to those of skill in the art.

5           Methods for DNA sequencing are well known in the art. Conventional enzymatic methods employ DNA polymerase, Klenow fragment, Sequenase (US Biochemical Corp, Cleveland, OH) or Taq polymerase to extend DNA chains from an oligonucleotide primer annealed to the DNA template of interest. Methods have  
10       been developed for the use of both single-stranded and double-stranded templates. The chain termination reaction products may be electrophoresed on urea/polyacrylamide gels and detected either by autoradiography (for radionucleotide labeled precursors) or by fluorescence (for fluorescent-labeled precursors). Recent  
15       improvements in mechanized reaction preparation, sequencing and analysis using the fluorescent detection method have permitted expansion in the number of sequences that can be determined per day using machines such as the Applied Biosystems 377 DNA Sequencers (Applied Biosystems, Foster City, CA).

          The reading frame of the nucleotide sequence can be ascertained by several types of analyses. First, reading frames contained within the coding sequence can  
20       be analyzed for the presence of start codon ATG and stop codons TGA, TAA or TAG. Typically, one reading frame will continue throughout the major portion of a cDNA sequence while other reading frames tend to contain numerous stop codons. In such cases, reading frame determination is straightforward. In other more difficult cases, further analysis is required.

25           Algorithms have been created to analyze the occurrence of individual nucleotide bases at each putative codon triplet. See, for example J.W. Fickett, Nuc Acids Res 10:5303 (1982). Coding DNA for particular organisms (bacteria, plants and animals) tends to contain certain nucleotides within certain triplet periodicities, such as a significant preference for pyrimidines in the third codon position. These  
30       preferences have been incorporated into widely available software which can be used to determine coding potential (and frame) of a given stretch of DNA. The algorithm-derived information combined with start/stop codon information can be used to determine proper frame with a high degree of certainty. This, in turn, readily permits cloning of the sequence in the correct reading frame into appropriate  
35       expression vectors.

[illegible][illegible][illegible][illegible]

priming and transcription, as well as a number of unique restriction sites, including EcoRI, for cloning. The vector can be transfected into an appropriate host strain of E. coli.

Induction of the isolated bacterial strain with isopropylthiogalactoside (IPTG) using standard methods will produce a fusion protein which contains the first seven residues of beta-galactosidase, about 15 residues of linker and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is one chance in three that the included cDNA will lie in the correct frame for proper translation. If the cDNA is not in the proper reading frame, the correct frame can be obtained by deletion or insertion of an appropriate number of bases by well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or oligonucleotide linker inclusion.

The cDNA can be shuttled into other vectors known to be useful for expression of protein in specific hosts. Oligonucleotide primers, containing cloning sites and segments of DNA sufficient to hybridize to stretches at both ends of the target cDNA, can be synthesized chemically by standard methods. These primers can then be used to amplify the desired gene segments by PCR. The resulting new gene segments can be digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments can be produced by digestion of the cDNA with appropriate restriction enzymes and filling in the missing gene segments with chemically synthesized oligonucleotides. Segments of the coding sequence from more than one gene can be ligated together and cloned in appropriate vectors to optimize expression of recombinant sequence.

Suitable expression hosts for such chimeric molecules include but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human embryonic kidney (HEK) 293 cells, insect cells such as Sf9 cells, yeast cells such as Saccharomyces cerevisiae and bacteria such as E. coli. For each of these cell systems, a useful expression vector may also include an origin of replication to allow propagation in bacteria and a selectable marker such as the beta-lactamase antibiotic resistance gene to allow selection in bacteria. In addition, the vectors may include a second selectable marker, such as the neomycin phosphotransferase gene, to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts may require the addition of 3' poly A tail if the sequence of interest lacks poly A.



Additionally, the vector may contain promoters or enhancers which increase gene expression. Such promoters are host specific and include, but are not limited to, MMTV, SV40, or metallothionine promoters for CHO cells; trp, lac, tac or T7 promoters for bacterial hosts; or alpha factor, alcohol oxidase or PGH promoters for yeast. Adenoviral vectors with or without transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to drive protein expression in mammalian cell lines. Once homogeneous cultures of recombinant cells are obtained, large quantities of recombinantly produced protein can be recovered from the conditioned medium and analyzed using chromatographic methods well known in the art. An alternative method for the production of large amounts of secreted protein involves the transfection of mammalian embryos and the recovery of the recombinant protein from milk produced by transgenic cows, goats, sheep, etc. Polypeptides and closely related molecules may be expressed recombinantly in such a way as to facilitate protein purification. One approach involves expression of a chimeric protein which includes one or more additional polypeptide domains not naturally present on human polypeptides. Such purification-facilitating domains include, but are not limited to, metal-chelating peptides such as histidine-tryptophan domains that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle, WA). The inclusion of a cleavable linker sequence such as Factor XA or enterokinase from Invitrogen (San Diego, CA) between the polypeptide sequence and the purification domain may be useful for recovering the polypeptide.

#### Immunoassays

BS106 polypeptides, including fragments, derivatives, and analogs thereof, or cells expressing such polypeptides, can be utilized in a variety of assays, many of which are described herein, for the detection of antibodies to breast tissue. They also can be used as immunogens to produce antibodies. These antibodies can be, for example, polyclonal or monoclonal antibodies, chimeric, single chain and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

For example, antibodies generated against a polypeptide comprising a sequence of the present invention can be obtained by direct injection of the polypeptide into an animal or by administering the polypeptide to an animal such as a

mouse, rabbit, goat or human. A mouse, rabbit or goat is preferred. The polypeptide is selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof. The antibody so obtained then will  
5 bind the polypeptide itself. In this manner, even a sequence encoding only a fragment of the polypeptide can be used to generate antibodies that bind the native polypeptide. Such antibodies then can be used to isolate the polypeptide from test samples such as tissue suspected of containing that polypeptide. For preparation of monoclonal antibodies, any technique which provides antibodies produced by  
10 continuous cell line cultures can be used. Examples include the hybridoma technique as described by Kohler and Milstein, Nature 256:495-497 (1975), the trioma technique, the human B-cell hybridoma technique as described by Kozbor et al, Immun. Today 4:72 (1983) and the EBV-hybridoma technique to produce human monoclonal antibodies as described by Cole et al., in Monoclonal Antibodies and  
15 Cancer Therapy, Alan R. Liss, Inc, New York, NY, pp. 77-96 (1985). Techniques described for the production of single chain antibodies can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. See, for example, U.S. Patent No. 4,946,778, which is incorporated herein by reference.

Various assay formats may utilize the antibodies of the present invention,  
20 including "sandwich" immunoassays and probe assays. For example, the antibodies of the present invention, or fragments thereof, can be employed in various assay systems to determine the presence, if any, of BS106 antigen in a test sample. For example, in a first assay format, a polyclonal or monoclonal antibody or fragment thereof, or a combination of these antibodies, which has been coated on a solid  
25 phase, is contacted with a test sample, to form a first mixture. This first mixture is incubated for a time and under conditions sufficient to form antigen/antibody complexes. Then, an indicator reagent comprising a monoclonal or a polyclonal antibody or a fragment thereof, or a combination of these antibodies, to which a signal generating compound has been attached, is contacted with the  
30 antigen/antibody complexes to form a second mixture. This second mixture then is incubated for a time and under conditions sufficient to form antibody/antigen/antibody complexes. The presence of BS106 antigen in the test sample and captured on the solid phase, if any, is determined by detecting the measurable signal generated by the signal generating compound. The amount of  
35 BS106 antigen present in the test sample is proportional to the signal generated.

1 In an alternative assay format, a mixture is formed by contacting: (1) a  
2 polyclonal antibody, monoclonal antibody, or fragment thereof, which specifically  
3 binds to BS106 antigen, or a combination of such antibodies bound to a solid  
4 support; (2) the test sample; and (3) an indicator reagent comprising a monoclonal  
5 antibody, polyclonal antibody, or fragment thereof, which specifically binds to a  
6 different BS106 antigen (or a combination of these antibodies) to which a signal  
7 generating compound is attached. This mixture is incubated for a time and under  
8 conditions sufficient to form antibody/antigen/antibody complexes. The presence, if  
9 any, of BS106 antigen present in the test sample and captured on the solid phase is  
10 determined by detecting the measurable signal generated by the signal generating  
11 compound. The amount of BS106 antigen present in the test sample is proportional  
12 to the signal generated.

13 In another assay format, one or a combination of at least two monoclonal  
14 antibodies of the invention can be employed as a competitive probe for the detection  
15 of antibodies to BS106 antigen. For example, BS106 polypeptides such as the  
16 recombinant antigens disclosed herein, either alone or in combination, are coated on  
17 a solid phase. A test sample suspected of containing antibody to BS106 antigen then  
18 is incubated with an indicator reagent comprising a signal generating compound and  
19 at least one monoclonal antibody of the invention for a time and under conditions  
20 sufficient to form antigen/antibody complexes of either the test sample and indicator  
21 reagent bound to the solid phase or the indicator reagent bound to the solid phase.  
22 The reduction in binding of the monoclonal antibody to the solid phase can be  
23 quantitatively measured.

24 In yet another detection method, each of the monoclonal or polyclonal  
25 antibodies of the present invention can be employed in the detection of BS106  
26 antigens in tissue sections, as well as in cells, by immunohistochemical analysis.  
27 Cytochemical analysis wherein these antibodies are labeled directly (with, for  
28 example, fluorescein, colloidal gold, horseradish peroxidase, alkaline phosphatase,  
29 etc.) or are labeled by using secondary labeled anti-species antibodies (with various  
30 labels as exemplified herein) to track the histopathology of disease also are within the  
31 scope of the present invention.

32 In addition, these monoclonal antibodies can be bound to matrices similar to  
33 CNBr-activated Sepharose and used for the affinity purification of specific BS106  
34 polypeptides from cell cultures or biological tissues such as to purify recombinant  
35 and native BS106 proteins.

The monoclonal antibodies of the invention also can be used for the generation of chimeric antibodies for therapeutic use, or other similar applications.

The monoclonal antibodies or fragments thereof can be provided individually to detect BS106 antigens. Combinations of the monoclonal antibodies (and  
5 fragments thereof) provided herein also may be used together as components in a mixture or "cocktail" of at least one BS106 antibody of the invention, along with antibodies which specifically bind to other BS106 regions, each antibody having different binding specificities. Thus, this cocktail can include the monoclonal antibodies of the invention which are directed to BS106 polypeptides disclosed  
10 herein and other monoclonal antibodies specific to other antigenic determinants of BS106 antigens or other related proteins.

The polyclonal antibody or fragment thereof which can be used in the assay formats should specifically bind to a BS106 polypeptide or other BS106 polypeptides additionally used in the assay. The polyclonal antibody used  
15 preferably is of mammalian origin such as, human, goat, rabbit or sheep polyclonal antibody which binds BS106 polypeptide. Most preferably, the polyclonal antibody is of rabbit origin. The polyclonal antibodies used in the assays can be used either alone or as a cocktail of polyclonal antibodies. Since the cocktails used in the assay formats are comprised of either monoclonal antibodies or polyclonal antibodies  
20 having different binding specificity to BS106 polypeptides, they are useful for the detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining the predisposition to, diseases and conditions of the breast such as breast cancer.

It is contemplated and within the scope of the present invention that BS106  
25 antigen may be detectable in assays by use of a recombinant antigen as well as by use of a synthetic peptide or purified peptide, which peptide comprises an amino acid sequence of BS106. The amino acid sequence of such a polypeptide is selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and  
30 fragments thereof. It also is within the scope of the present invention that different synthetic, recombinant or purified peptides, identifying different epitopes of BS106, can be used in combination in an assay for the detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining the predisposition to diseases and conditions of the breast such as breast cancer. In this  
35 case, all of these peptides can be coated onto one solid phase; or each separate

peptide may be coated onto separate solid phases, such as microparticles, and then combined to form a mixture of peptides which can be later used in assays.

Furthermore, it is contemplated that multiple peptides which define epitopes from different antigens may be used for the detection, diagnosis, staging, monitoring,  
5 prognosis, prevention or treatment of, or determining the predisposition to, diseases and conditions of the breast, such as breast cancer. Peptides coated on solid phases or labeled with detectable labels are then allowed to compete with those present in a patient sample (if any) for a limited amount of antibody. A reduction in binding of the synthetic, recombinant, or purified peptides to the antibody (or antibodies) is an  
10 indication of the presence of BS106 antigen in the patient sample. The presence of BS106 antigen indicates the presence of breast tissue disease, especially breast cancer, in the patient. Variations of assay formats are known to those of ordinary skill in the art and many are discussed herein below.

In another assay format, the presence of anti-BS106 antibody and/or BS106  
15 antigen can be detected in a simultaneous assay, as follows. A test sample is simultaneously contacted with a capture reagent of a first analyte, wherein said capture reagent comprises a first binding member specific for a first analyte attached to a solid phase and a capture reagent for a second analyte, wherein said capture reagent comprises a first binding member for a second analyte attached to a second  
20 solid phase, to thereby form a mixture. This mixture is incubated for a time and under conditions sufficient to form capture reagent/first analyte and capture reagent/second analyte complexes. These so-formed complexes then are contacted with an indicator reagent comprising a member of a binding pair specific for the first analyte labeled with a signal generating compound and an indicator reagent  
25 comprising a member of a binding pair specific for the second analyte labeled with a signal generating compound to form a second mixture. This second mixture is incubated for a time and under conditions sufficient to form capture reagent/first analyte/indicator reagent complexes and capture reagent/second analyte/indicator reagent complexes. The presence of one or more analytes is determined by detecting  
30 a signal generated in connection with the complexes formed on either or both solid phases as an indication of the presence of one or more analytes in the test sample. In this assay format, recombinant antigens derived from the expression systems disclosed herein may be utilized, as well as monoclonal antibodies produced from the proteins derived from the expression systems as disclosed herein. For example,

in this assay system, BS106 antigen can be the first analyte. Such assay systems are described in greater detail in EP Publication No. 0473065.

In yet other assay formats, the polypeptides disclosed herein may be utilized to detect the presence of antibody against BS106 antigen in test samples. For  
5 example, a test sample is incubated with a solid phase to which at least one polypeptide such as a recombinant protein or synthetic peptide has been attached. The polypeptide is selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof. These are reacted for a time and  
10 under conditions sufficient to form antigen/antibody complexes. Following incubation, the antigen/antibody complex is detected. Indicator reagents may be used to facilitate detection, depending upon the assay system chosen. In another assay format, a test sample is contacted with a solid phase to which a recombinant protein produced as described herein is attached, and also is contacted with a  
15 monoclonal or polyclonal antibody specific for the protein, which preferably has been labeled with an indicator reagent. After incubation for a time and under conditions sufficient for antibody/antigen complexes to form, the solid phase is separated from the free phase, and the label is detected in either the solid or free phase as an indication of the presence of antibody against BS106 antigen. Other  
20 assay formats utilizing the recombinant antigens disclosed herein are contemplated. These include contacting a test sample with a solid phase to which at least one antigen from a first source has been attached, incubating the solid phase and test sample for a time and under conditions sufficient to form antigen/antibody complexes, and then contacting the solid phase with a labeled antigen, which antigen  
25 is derived from a second source different from the first source. For example, a recombinant protein derived from a first source such as E. coli is used as a capture antigen on a solid phase, a test sample is added to the so-prepared solid phase, and following standard incubation and washing steps as deemed or required, a recombinant protein derived from a different source (i.e., non-E. coli) is utilized as a  
30 part of an indicator reagent which subsequently is detected. Likewise, combinations of a recombinant antigen on a solid phase and synthetic peptide in the indicator phase also are possible. Any assay format which utilizes an antigen specific for BS106 produced or derived from a first source as the capture antigen and an antigen specific for BS106 from a different second source is contemplated. Thus, various  
35 combinations of recombinant antigens, as well as the use of synthetic peptides,

purified proteins and the like, are within the scope of this invention. Assays such as this and others are described in U.S. Patent No. 5,254,458, which enjoys common ownership and is incorporated herein by reference.

Other embodiments which utilize various other solid phases also are contemplated and are within the scope of this invention. For example, ion capture procedures for immobilizing an immobilizable reaction complex with a negatively charged polymer (described in EP publication 0326100 and EP publication No. 0406473), can be employed according to the present invention to effect a fast solution-phase immunochemical reaction. An immobilizable immune complex is separated from the rest of the reaction mixture by ionic interactions between the negatively charged poly-anion/immune complex and the previously treated, positively charged porous matrix and detected by using various signal generating systems previously described, including those described in chemiluminescent signal measurements as described in EPO Publication No. 0 273,115.

Also, the methods of the present invention can be adapted for use in systems which utilize microparticle technology including automated and semi-automated systems wherein the solid phase comprises a microparticle (magnetic or non-magnetic). Such systems include those described in, for example, published EPO applications Nos. EP 0 425 633 and EP 0 424 634, respectively.

The use of scanning probe microscopy (SPM) for immunoassays also is a technology to which the monoclonal antibodies of the present invention are easily adaptable. In scanning probe microscopy, particularly in atomic force microscopy, the capture phase, for example, at least one of the monoclonal antibodies of the invention, is adhered to a solid phase and a scanning probe microscope is utilized to detect antigen/antibody complexes which may be present on the surface of the solid phase. The use of scanning tunneling microscopy eliminates the need for labels which normally must be utilized in many immunoassay systems to detect antigen/antibody complexes. The use of SPM to monitor specific binding reactions can occur in many ways. In one embodiment, one member of a specific binding partner (analyte specific substance which is the monoclonal antibody of the invention) is attached to a surface suitable for scanning. The attachment of the analyte specific substance may be by adsorption to a test piece which comprises a solid phase of a plastic or metal surface, following methods known to those of ordinary skill in the art. Or, covalent attachment of a specific binding partner (analyte specific substance) to a test piece which test piece comprises a solid phase of

derivatized plastic, metal, silicon, or glass may be utilized. Covalent attachment methods are known to those skilled in the art and include a variety of means to irreversibly link specific binding partners to the test piece. If the test piece is silicon or glass, the surface must be activated prior to attaching the specific binding partner.

5 Also, polyelectrolyte interactions may be used to immobilize a specific binding partner on a surface of a test piece by using techniques and chemistries. The preferred method of attachment is by covalent means. Following attachment of a specific binding member, the surface may be further treated with materials such as serum, proteins, or other blocking agents to minimize non-specific binding. The  
10 surface also may be scanned either at the site of manufacture or point of use to verify its suitability for assay purposes. The scanning process is not anticipated to alter the specific binding properties of the test piece.

While the present invention discloses the preference for the use of solid phases, it is contemplated that the reagents such as antibodies, proteins and peptides  
15 of the present invention can be utilized in non-solid phase assay systems. These assay systems are known to those skilled in the art, and are considered to be within the scope of the present invention.

It is contemplated that the reagent employed for the assay can be provided in the form of a test kit with one or more containers such as vials or bottles, with each  
20 container containing a separate reagent such as a probe, primer, monoclonal antibody or a cocktail of monoclonal antibodies, or a polypeptide (e.g. recombinantly, synthetically produced or purified) employed in the assay. The polypeptide is selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and  
25 fragments thereof. Other components such as buffers, controls and the like, known to those of ordinary skill in art, may be included in such test kits. It also is contemplated to provide test kits which have means for collecting test samples comprising accessible body fluids, e.g., blood, urine, saliva and stool. Such tools useful for collection ("collection materials") include lancets and absorbent paper or  
30 cloth for collecting and stabilizing blood; swabs for collecting and stabilizing saliva; cups for collecting and stabilizing urine or stool samples. Collection materials, papers, cloths, swabs, cups and the like, may optionally be treated to avoid denaturation or irreversible adsorption of the sample. The collection materials also may be treated with or contain preservatives, stabilizers or antimicrobial agents to  
35 help maintain the integrity of the specimens. Test kits designed for the collection,



stabilization and preservation of test specimens obtained by surgery or needle biopsy are also useful. It is contemplated that all kits may be configured in two components which can be provided separately; one component for collection and transport of the specimen and the other component for the analysis of the specimen. The collection component, for example, can be provided to the open market user while the components for analysis can be provided to others such as laboratory personnel for determination of the presence, absence or amount of analyte. Further, kits for the collection, stabilization and preservation of test specimens may be configured for use by untrained personnel and may be available in the open market for use at home with subsequent transportation to a laboratory for analysis of the test sample.

E. coli bacteria (clone 1662885) has been deposited at the American Type Culture Collection (A.T.C.C.), 12301 Parklawn Drive, Rockville, Maryland 20852, as of 11/20/96, under the terms of the Budapest Treaty and will be maintained for a period of thirty (30) years from the date of deposit, or for five (5) years after the last request for the deposit, or for the enforceable period of the U.S. patent, whichever is longer. The deposit and any other deposited material described herein are provided for convenience only, and are not required to practice the present invention in view of the teachings provided herein. The cDNA sequence in all of the deposited material is incorporated herein by reference. Clone 1662885 was accorded A.T.C.C. Deposit No. 98256.

The present invention will now be described by way of examples, which are meant to illustrate, but not to limit, the scope of the present invention.

#### EXAMPLES

##### Example 1: Identification of Breast Tissue Library BS106 Gene-Specific Clones

A. Library Comparison of Expressed Sequence Tags (ESTs) or Transcript Images. Partial sequences of cDNA clone inserts, so-called "expressed sequence tags" (ESTs), were derived from cDNA libraries made from breast tumor tissues, breast non-tumor tissues and numerous other tissues, both tumor and non-tumor and entered into a database (LIFESEQ™ database, available from Incyte Pharmaceuticals, Palo Alto, CA) as gene transcript images. See International Publication No. WO 95/20681. (A transcript image is a listing of the number of ESTs for each of the represented genes in a given tissue library. ESTs sharing regions of mutual sequence overlap are classified into clusters. A cluster is assigned a clone number from a representative 5' EST. Often, a cluster of interest can be

extended by comparing its consensus sequence with sequences of other ESTs which did not meet the criteria for automated clustering. The alignment of all available clusters and single ESTs represent a contig from which a consensus sequence is derived.) The transcript images then were evaluated to identify EST clusters that  
5 were representative primarily of the breast tissue libraries. These target clusters then were ranked according to their abundance (occurrence of EST members) in the target libraries and their absence from background libraries. Higher abundance clusters with low background occurrence were given higher study priority. Overlapping clones 1662885 (SEQUENCE ID NO 1), 893988 (SEQUENCE ID NO 2), and  
10 1209814 (SEQUENCE ID NO 3) were identified for further study. These represented the minimum number of clones that were needed to form the BS106 contig and from which the consensus sequence provided herein (SEQUENCE ID NO 4) was derived.

B. Generation of a Consensus Sequence. The nucleotide sequences of EST  
15 clones, 1662885 (SEQUENCE ID NO 1), 893988 (SEQUENCE ID NO 2), and 1209814 (SEQUENCE ID NO 3), were entered in the Sequencher™ Program (available from Gene Codes Corporation, Ann Arbor, MI, in order to generate a nucleotide alignment (contig map) and then generate their consensus sequence (SEQUENCE ID NO 4). FIGURE 1 shows the nucleotide sequence alignment of  
20 these clones and their resultant nucleotide consensus sequence (SEQUENCE ID NO 4). FIGURE 2 presents the contig map depicting the clones 1662885 (SEQUENCE ID NO 1), 893988 (SEQUENCE ID NO 2), and 1209814 (SEQUENCE ID NO 3) forming overlapping regions of the BS106 gene and the resultant consensus nucleotide sequence (SEQUENCE ID NO 4) of these clones in a graphic display.  
25 Following this, a three-frame translation was performed on the consensus sequence (SEQUENCE ID NO 4). The first forward frame was found to have an open reading frame encoding a 90 residue amino acid sequence, which is presented as SEQUENCE ID NO 16. FIGURE 1 shows a G/T polymorphism at position 54 in the consensus sequence (indicated as "K"); the respective two nucleotides were  
30 noted with approximately equal frequency in the LIFESEQ™ database. It is believed that the polymorphism at position 54 which lies 10 nucleotides upstream of a methionine residue (the potential translational start site) may be involved in regulating the level of expression of this protein. FIGURE 1 also shows a C/T polymorphism at position 312 in the consensus sequence (indicated as "Y"). The  
35 ratio of C's to T's noted in the LIFESEQ™ database at position 312 was 3:1.

SEQUENCE ID NO 4

C. Specificity of Expression of ESTs Corresponding to Consensus

Sequence. The consensus sequence, generated in section B, supra, was compared to the entire updated LIFESEQ™ database (September 1997) using the BLAST search tool. ESTs corresponding to the consensus sequence were found in 85.7% (24 of 28) of breast libraries and 0.2% (1 of 455) of non-breast libraries. Therefore, the consensus sequence, or fragments thereof, were found more than 389 times more often in breast than non-breast tissues.

Example 2: Sequencing of BS106 EST-Specific Clones

The DNA sequence of clone 1662885 (SEQUENCE ID NO 5) which comprises the 5'-most EST of the BS106 gene contig was determined using dideoxy termination sequencing with dye terminators following known methods. (F. Sanger et al., PNAS U.S.A. 74:5463 (1977)). An A/G polymorphism at position 543 in SEQUENCE ID NO 5 is indicated as "R."

Because the pINCY vector (Life Technologies, Gaithersburg, MD) contains universal priming sites just adjacent to the 3' and 5' ligation junctions of the inserts, approximately 300 bases of the insert were sequenced in both directions using universal primers, SEQUENCE ID NO 8 and SEQUENCE ID NO 9 (New England Biolabs, Beverly, MA and Applied Biosystems Inc, Foster City, CA, respectively).

The sequencing reactions were run on a polyacrylamide denaturing gel, and the sequences were determined by an Applied Biosystems 377 Sequencer (available from Applied Biosystems, Foster City, CA) or other sequencing apparatus. Additional sequencing primers, BS106.F1 and BS106.R1 (SEQUENCE ID NO 10 and SEQUENCE ID NO 11, respectively) were designed from sequence determined by the initial sequencing reactions near the 3'-ends of the two DNA strands. These primers then were used to determine the remaining DNA sequence of the cloned insert from each DNA strand, as previously described.

Example 3: Nucleic Acid Preparation

A. RNA Extraction from Tissue. Total RNA was isolated from solid breast tissues or cells and from non-breast tissues. Various methods were utilized, including but not limited to the lithium chloride/urea technique, known and described in the art (Kato et al., J. Virol. 61:2182-2191, (1987)), Ultraspec™ (Biotecx Laboratories, Inc., Houston Texas), and TRIzol™ (Life Technologies, Inc., Gaithersburg, MD).

For northern blot analysis, the tissue was placed in a sterile conical tube on ice and 10-15 volumes of 3 M LiCl, 6 M urea, 5 mM EDTA, 0.1 M  $\beta$ -mercaptoethanol, 50 mM Tris-HCl (pH 7.5) were added. The tissue was homogenized with a Polytron® homogenizer (Brinkman Instruments, Inc.,  
5 Westbury, NY) for 30-50 sec on ice. The solution was transferred to a 15 ml plastic centrifuge tube and placed overnight at -20°C. The tube was centrifuged for 90 min at 9,000 x g at 0-4°C, and the supernatant was immediately decanted. Then, 10 ml of 3 M LiCl were added, the tube was vortexed for 5 sec and centrifuged for 45 min at 11,000 x g at 0-4°C. These decanting, resuspension in LiCl, and centrifugation  
10 steps were repeated. The final pellet was air dried and resuspended in 2 ml of 1 mM EDTA, 0.5% SDS, 10 mM Tris (pH 7.5). Then, a 20  $\mu$ l volume of Proteinase K (20 mg/ml) was added, and the resulting solution was incubated for 30 min at 37°C with occasional mixing. One-tenth volume (0.22-0.25 ml) of 3 M NaCl was added and the solution was vortexed before transfer into another tube which contained 2 ml  
15 of phenol/chloroform/isoamyl alcohol (PCI). The tube was vortexed for 1-3 sec and centrifuged for 20 min at 3,000 x g at 10°C. The PCI extraction was repeated twice more, followed by two similar extractions with chloroform/isoamyl alcohol. The final aqueous solution was transferred to a pre-chilled 15 ml corex glass tube containing 6 ml of 100% absolute ethanol, the tube was covered with parafilm and  
20 placed at -20°C overnight. The tube was centrifuged for 30 min at 10,000 x g at 0-4°C, and the ethanol supernatant was decanted immediately. The RNA pellet was washed four times with 10 ml of 75% ice-cold ethanol, followed each time by centrifugation at 10,000 x g for 10 min. The final pellet was air dried for 15 min at room temperature. The RNA was suspended in 0.5 ml of 10 mM Tris (pH 7.6), 1  
25 mM EDTA, and its concentration was determined spectrophotometrically. RNA samples were aliquoted and stored at -70°C as ethanol precipitates.

The quality of the RNA was determined by agarose gel electrophoresis (see Example 5) and staining with 0.5  $\mu$ g/ml ethidium bromide for one hour. RNA samples that did not contain intact 28S/18S rRNAs were excluded from the study.

30 Alternatively, for RT-PCR analysis, 1 ml of Ultraspec RNA reagent was added to 120 mg of pulverized tissue in a 2.0 ml polypropylene microfuge tube, homogenized with a Polytron® homogenizer (Brinkman Instruments, Inc., Westbury, NY) for 50 sec, and left on ice for 5 min. Then, 0.2 ml of chloroform was added to each sample, followed by vortexing for 15 sec. The sample was left in  
35 ice for another 5 min, followed by centrifugation at 12,000 x g for 15 min at 4°C.

The upper layer was collected and transferred to another RNase-free 2.0 ml microfuge tube. An equal volume of isopropanol was added to each sample, and the solution was placed on ice for 10 min. The sample was centrifuged at 12,000 x g for 10 min at 4°C, and the supernatant was discarded. The remaining pellet was  
5 washed twice with cold 75% ethanol, resuspended by vortexing, and the resuspended material was then re-pelleted by centrifugation at 7500 x g for 5 min at 4°C. Finally, the RNA pellet was dried in a speedvac for at least 5 min and reconstituted in RNase-free water.

B. RNA Extraction from Blood Mononuclear Cells. Mononuclear cells are  
10 isolated from blood samples from patients by centrifugation using Ficoll-Hypaque as follows. A 10 ml volume of whole blood is mixed with an equal volume of RPMI Medium (Life Technologies, Gaithersburg, MD). This mixture is then underlayered with 10 ml of Ficoll-Hypaque (Pharmacia, Piscataway, NJ) and centrifuged for 30 minutes at 200 x g. The buffy coat containing the mononuclear cells is removed,  
15 diluted to 50 ml with Dulbecco's PBS (Life Technologies, Gaithersburg, MD) and the mixture centrifuged for 10 minutes at 200 x g. After two washes, the resulting pellet is resuspended in Dulbecco's PBS to a final volume of 1 ml.

RNA is prepared from the isolated mononuclear cells as described by N. Kato et al., supra. Briefly, the pelleted mononuclear cells are brought to a final of 1  
20 ml volume and then are resuspended in 250 µL of PBS and mixed with 2.5 ml of 3M LiCl, 6M urea, 5mM EDTA, 0.1M 2-mercaptoethanol, 50mM Tris-HCl (pH 7.5). The resulting mixture is homogenized and incubated at -20°C overnight. The homogenate is spun at 8,000 RPM in a Beckman J2-21M rotor for 90 minutes at 0-4°C. The pellet is resuspended in 10 ml 3M LiCl by vortexing and then spun at  
25 10,000 RPM in a Beckman J2-21M rotor centrifuge for 45 minutes at 0-4°C. The resuspending and pelleting steps then are repeated. The pellet is resuspended in 2 ml of 1 mM EDTA, 0.5% SDS, 10 mM Tris (pH 7.5) and 400 µg Proteinase K with vortexing and then it is incubated at 37°C for 30 minutes with shaking. One tenth volume of 3M NaCl then is added and the vortexed mixture. Proteins are removed  
30 by two cycles of extraction with phenol/ chloroform/ isoamyl alcohol followed by one extraction with chloroform/ isoamyl alcohol. RNA is precipitated by the addition of 6 ml of ethanol followed by overnight incubation at -20°C. After the precipitated RNA is collected by centrifugation, the pellet is washed 4 times in 75% ethanol. The pelleted RNA is then dissolved in 1mM EDTA, 10mM Tris-HCl (pH  
35 7.5).

Non-breast tissues are used as negative controls. The mRNA can be further purified from total RNA by using commercially available kits such as oligo dT cellulose spin columns (RediCol™ from Pharmacia, Uppsala, Sweden) for the isolation of poly-adenylated RNA. Total or mRNA can be dissolved in lysis buffer  
5 (5M guanidine thiocyanate, 0.1M EDTA, pH 7.0) for analysis in the ribonuclease protection assay.

C. RNA Extraction from polysomes. Tissue is minced in saline at 4°C and mixed with 2.5 volumes of 0.8 M sucrose in a TK<sub>150</sub>M (150 mM KCl, 5 mM MgCl<sub>2</sub>, 50 mM Tris-HCl, pH 7.4) solution containing 6 mM 2-mercaptoethanol.  
10 The tissue is homogenized in a Teflon-glass Potter homogenizer with five strokes at 100-200 rpm followed by six strokes in a Dounce homogenizer, as described by B. Mechler, Methods in Enzymology 152:241-248 (1987). The homogenate then is centrifuged at 12,000 x g for 15 min at 4°C to sediment the nuclei. The polysomes are isolated by mixing 2 ml of the supernatant with 6 ml of 2.5 M sucrose in TK<sub>150</sub>M and layering this mixture over 4 ml of 2.5 M sucrose in TK<sub>150</sub>M in a 38 ml  
15 polyallomer tube. Two additional sucrose TK<sub>150</sub>M solutions are successively layered onto the extract fraction; a first layer of 13 ml 2.05 M sucrose followed by a second layer of 6 ml of 1.3 M sucrose. The polysomes are isolated by centrifuging the gradient at 90,000 x g for 5 h at 4°C. The fraction then is taken from the 1.3 M  
20 sucrose/2.05 M sucrose interface with a siliconized pasteur pipette and diluted in an equal volume of TE (10 mM Tris-HCl, pH 7.4, 1 mM EDTA). An equal volume of 90°C SDS buffer (1% SDS, 200 mM NaCl, 20 mM Tris-HCl, pH 7.4) is added and the solution is incubated in a boiling water bath for 2 min. Proteins next are digested with a Proteinase-K digestion (50 mg/ml) for 15 min at 37°C. The mRNA is  
25 purified with 3 equal volumes of phenol-chloroform extractions followed by precipitation with 0.1 volume of 2 M sodium acetate (pH 5.2) and 2 volumes of 100% ethanol at -20°C overnight. The precipitated RNA is recovered by centrifugation at 12,000 x g for 10 min at 4°C. The RNA is dried and resuspended in TE (pH 7.4) or distilled water. The resuspended RNA then can be used in a slot  
30 blot or dot blot hybridization assay to check for the presence of BS106 mRNA (see Example 6).

The quality of nucleic acid and proteins is dependent on the method of preparation used. Each sample may require a different preparation technique to maximize isolation efficiency of the target molecule. These preparation techniques  
35 are within the skill of the ordinary artisan.

Example 4: Ribonuclease Protection Assay

A. Synthesis of Labeled Complementary RNA (cRNA) Hybridization Probe and Unlabeled Sense Strand. A pINCY plasmid containing the BS106 gene cDNA sequence insert (clone 1662885), flanked by opposed SP6 and T7 polymerase promoters, was purified using Qiagen Plasmid Purification Kit (Qiagen, Chatsworth, CA). Then, 10 µg of the plasmid were cut with 10 U DdeI restriction enzyme for 1 h at 37°C. The cut plasmid was purified using QIAprep kits (Qiagen, Chatsworth, CA) and used for the synthesis of antisense transcript labeled with 6.3 µM (alpha<sup>32</sup>P) UTP (800 Ci/mmol)(Amersham Life Sciences, Inc. Arlington Heights, IL) from the SP6 promoter using the Riboprobe<sup>®</sup> in vitro Transcription System (Promega Corporation, Madison, WI), as described by the supplier's instructions. To generate the sense strand, 10 µg of the purified plasmid were cut with restriction enzymes 10 U XbaI and 10 U NotI, and transcribed as above from the T7 promoter. Both sense and antisense strands were isolated by spin column chromatography. Unlabeled sense strand was quantitated by UV absorption at 260 nm.

B. Hybridization of Labeled Probe to Target. Frozen tissue was pulverized to powder under liquid nitrogen and 100-500 mg were dissolved in 1 ml of lysis buffer as available as a component of the DirectProtect<sup>™</sup> Lysate RNase Protection kit (Ambion, Inc., Austin, TX). Further dissolution was achieved using a tissue homogenizer. In addition, a dilution series of a known amount of sense strand in mouse liver lysate was made for use as a positive control. Finally, 45 µl of solubilized tissue or diluted sense strand was mixed directly with 1 x 10<sup>5</sup> cpm of radioactively labeled probe (1.5 x 10<sup>9</sup> cpm/µg) in 5 µl of lysis buffer. Hybridization was allowed to proceed overnight at 37°C.

C. RNase Digestion. RNA that was not hybridized to probe was removed from the reaction as per the Direct Protect<sup>™</sup> protocol using a solution of RNase A and RNase T1 for 30 min at 37°C, followed by removal of RNase by Proteinase-K digestion in the presence of sodium sarcosyl. Hybridized fragments protected from digestion were then precipitated by the addition of an equal volume of isopropanol and placed at -70°C for 3 h. The precipitates were collected by centrifugation at 12,000 x g for 20 min.

D. Fragment Analysis. The precipitates were dissolved in denaturing gel loading dye (80% formamide, 10 mM EDTA (pH 8.0), 1 mg/ml xylene cyanol, 1

mg/ml bromophenol blue), heat denatured, and electrophoresed in 6% polyacrylamide TBE, 8 M urea denaturing gels. The gels were imaged and analyzed using the STORM™ storage phosphor autoradiography system (Molecular Dynamics, Sunnyvale, CA). Quantitation of protected fragment bands, expressed in femtograms (fg), was achieved by comparing the peak areas obtained from the test samples to those from the known dilutions of the positive control sense strand (see Section B, supra). In addition, the concentration of DNA in the lysate was assayed to estimate the number of cells in the test sample lysates. The results are expressed in molecules of BS106 RNA/cell and as a image rating score (Table 1). High level expression of mRNA corresponding to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof, indicated the presence of BS106 mRNA(s), suggesting a diagnosis of a breast tissue disease or condition, such as breast cancer.

Table 1

| Tissue           | ID Number | BS106 RNA/cell | Score* |
|------------------|-----------|----------------|--------|
| Normal Breast    | C157      | 10             | +      |
|                  | C007G     | 7              | +      |
|                  | C027R     | 8              | +      |
|                  | C016R     | 2              | +      |
|                  | C135R     | 0.2            | +      |
| Malignant Breast | C011G     | >46            | 3+     |
|                  | C023G     | 0.3            | +      |
|                  | C012G     | 0              | -      |
|                  | C033R     | >87            | 3+     |
|                  | C030      | 0              | -      |
| Normal Lung      | C005R     | 0              | -      |
| Malignant Lung   | C037G     | 0              | -      |
| Normal Colon     | C027G     | 0              | -      |

\*Samples with no detectable protected fragment were scored "-"; samples with detectable protected fragment (the fg values of which were within the std curve) were scored "+"; samples with detectable protected fragment, the fg values of which were 2 to 10 fold above the std curve, were scored "2+"; samples with detectable protected fragment, the fg values of which were 10 fold or more above the std curve, were scored "3+".

#### Example 5: Northern Blotting

The northern blot technique was used to identify a specific size RNA species in a complex population of RNA using agarose gel electrophoresis



and nucleic acid hybridization. Briefly, 5 µg of total RNA (see Example 3) was incubated in 15 µl of a solution containing 40 mM morpholinopropanesulfonic acid (MOPS) (pH 7.0), 10 mM sodium acetate, 1 mM EDTA, 2.2 M formaldehyde, 50% v/v formamide for 15 min at 65°C. The denatured RNA was mixed with 2 liters of loading buffer (50% glycerol, 1 mM EDTA, 0.4% bromophenol blue, 0.4% xylene cyanol) and loaded on a denaturing 1.0% agarose gel containing 40 mM MOPS (pH 7.0), 10 mM sodium acetate, 1 mM EDTA and 2.2 M formaldehyde. The gel was electrophoresed at 60 V for 1.5 h and rinsed in RNase free water. Gels were stained with 0.5 g/ml of ethidium bromide in RNase free water and illuminated with UV light to visualize ribosomal RNA bands. RNA was transferred from the gel onto nylon membranes (Brightstar-Plus, Ambion, Inc., Austin, TX) for 1.5 hours using the downward alkaline capillary transfer method (Chomczynski, Anal. Biochem. 201:134-139, 1992). The filter was rinsed with 1X SSC, and RNA was crosslinked to the filter using a Stratalinker (Stratagene, Inc., La Jolla, CA) on the autocrosslinking mode and dried for 15 min. The membrane was then placed into a hybridization tube containing 20 ml of preheated prehybridization solution (5X SSC, 50% formamide, 5X Denhardt's solution, 100 g/ml denatured salmon sperm DNA) and incubated in a 42°C hybridization oven for at least 3 hr. While the blot was prehybridizing, a 32P-labeled random-primed probe was generated using the BS106 insert fragment (obtained by digesting clone 1662885 with XbaI and NotI) using Random Primer DNA Labeling System (Life Technologies, Inc., Gaithersburg, MD) according to the manufacturer's instructions. Half of the probe was boiled for 10 min, quick chilled on ice and added to the hybridization tube. Hybridization was carried out at 42°C for at least 12 hr. The hybridization solution was discarded and the filter was washed twice in 30 ml of 3X SSC, 0.1% SDS at 42°C for 15 min, followed by two washes with 0.3X SSC, 0.1% SDS at 60°C for 15 min. The filter was wrapped in saran wrap, exposed to Kodak XAR-Omat film for 24-96 hr, and the film was developed for analysis. Results of the analysis of RNA quality using an ethidium bromide stained agarose gel and the corresponding northern blot using BS106 probe hybridized to RNAs from breast tissues and prostate tissues, and from normal breast and breast cancer tissues in Figures 3A and 3B, respectively. The positions of RNA size standards (in Kb) are shown to the left of each panel. As

shown in Figure 3A, the BS106 probe hybridized to an RNA band of approximately 0.7 Kb in 3 of 3 normal breast tissues, and in 1 of 3 prostate cancer tissues; but, an RNA band was not detected in 3 normal prostate tissues or in a prostate or breast cancer cell line. Figure 3B shows that the BS106 probe hybridized to an RNA of approximately 0.7 Kb in five of 6 normal breast RNA samples and 2 of 6 breast cancer RNA samples.

#### Example 6: Dot Blot/Slot Blot

Dot and slot blot assays are quick methods to evaluate the presence of a specific nucleic acid sequence in a complex mix of nucleic acid. To perform such assays, up to 50 µg of RNA is mixed in 50 µl of 50% formamide, 7% formaldehyde, 1X SSC, incubated 15 min at 68°C, and then cooled on ice. Then, 100 µl of 20X SSC is added to the RNA mixture and loaded under vacuum onto a manifold apparatus that has a prepared nitrocellulose or nylon membrane. The membrane is soaked in water, 20X SSC for 1 hour, placed on two sheets of 20X SSC prewet Whatman #3 filter paper, and loaded into a slot blot or dot blot vacuum manifold apparatus. The slot blot is analyzed with probes prepared and labeled as described in Example 4, supra. Detection of mRNA corresponding to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof, is an indication of the presence of BS106, suggesting a diagnosis of a breast tissue disease or condition, such as breast cancer.

Other methods and buffers which can be utilized in the methods described in Examples 5 and 6, but not specifically detailed herein, are known in the art and are described in J. Sambrook et al, supra which is incorporated herein by reference.

#### Example 7: In Situ Hybridization

This method is useful to directly detect specific target nucleic acid sequences in cells using detectable nucleic acid hybridization probes.

Tissues are prepared with cross-linking fixative agents such as paraformaldehyde or glutaraldehyde for maximum cellular RNA retention. See, L. Angerer et al., Methods in Cell Biol. 35:37-71 (1991). Briefly, the tissue is placed in greater than 5 volumes of 1% glutaraldehyde in 50 mM sodium phosphate, pH 7.5 at 4°C for 30 min. The solution is changed with fresh glutaraldehyde solution (1% glutaraldehyde in 50mM sodium phosphate, pH 7.5) for a further 30 min

fixing. The fixing solution should have an osmolality of approximately 0.375% NaCl. The tissue is washed once in isotonic NaCl to remove the phosphate.

The fixed tissues then are embedded in paraffin as follows. The tissue is dehydrated through a series of ethanol concentrations for 15 min each: 50% (twice),  
5 70% (twice), 85%, 90% and then 100% (twice). Next, the tissue is soaked in two changes of xylene for 20 min each at room temperature. The tissue is then soaked in two changes of a 1:1 mixture of xylene and paraffin for 20 min each at 60°C; and then in three final changes of paraffin for 15 min each.

Next, the tissue is cut in 5 µm sections using a standard microtome and  
10 placed on a slide previously treated with a tissue adhesive such as 3-aminopropyltriethoxysilane.

Paraffin is removed from the tissue by two 10 min xylene soaks and rehydrated in a series of ethanol concentrations: 99% twice, 95%, 85%, 70%, 50%, 30%, and then distilled water twice. The sections are pre-treated with 0.2 M HCl for  
15 10 min and permeabilized with 2 µg/ml Proteinase-K at 37°C for 15 min.

Labeled riboprobes transcribed from the BS106 gene plasmid (see Example 4) are hybridized to the prepared tissue sections and incubated overnight at 56°C in 3X standard saline extract and 50% formamide. Excess probe is removed by washing in 2X standard saline citrate and 50% formamide followed by digestion  
20 with 100 µg/ml RNase A at 37°C for 30 min. Fluorescence probe is visualized by illumination with ultraviolet (UV) light under a microscope. Fluorescence in the cytoplasm is indicative of BS106 mRNA. Alternatively, the sections can be visualized by autoradiography.

25 Example 8: Reverse Transcription PCR

A. One Step RT-PCR Assay. Target-specific primers are designed to detect the above-described target sequences by reverse transcription PCR using methods known in the art. One step RT-PCR is a sequential procedure that performs both RT and PCR in a single reaction mixture. The procedure is performed in a 200 µl  
30 reaction mixture containing 50 mM (N,N,-bis[2-Hydroxyethyl]glycine), pH 8.15, 81.7 mM KOAc, 33.33 mM KOH, 0.01 mg/ml bovine serum albumin, 0.1 mM ethylene diaminetetraacetic acid, 0.02 mg/ml NaN<sub>3</sub>, 8% w/v glycerol, 150 µM each of dNTP, 0.25 µM each primer, 5U rTth polymerase, 3.25 mM Mn(OAc)<sub>2</sub> and 5 µl of target RNA (see Example 3). Since RNA and the rTth polymerase enzyme are  
35 unstable in the presence of Mn(OAc)<sub>2</sub>, the Mn(OAc)<sub>2</sub> should be added just before

target addition. Optimal conditions for cDNA synthesis and thermal cycling readily can be determined by those skilled in the art. The reaction is incubated in a Perkin-Elmer Thermal Cycler 480. Optimal conditions for cDNA synthesis and thermal cycling can readily be determined by those skilled in the art. Conditions which may  
5 be found useful include cDNA synthesis at 60°-70°C for 15-45 min and 30-45 amplification cycles at 94°C, 1 min; 55°-70°C, 1 min; 72°C, 2 min. One step RT-PCR also may be performed by using a dual enzyme procedure with Taq polymerase and a reverse transcriptase enzyme, such as MMLV or AMV RT enzymes.

B. Traditional RT-PCR. A traditional two-step RT-PCR reaction was  
10 performed, as described by K.Q. Hu et al., *Virology* 181:721-726 (1991). Briefly, 0.5 µg of extracted mRNA (see Example 3) was reverse transcribed in a 20 µl reaction mixture containing 1X PCR II buffer (Perkin-Elmer), 5 mM MgCl<sub>2</sub>, 1 mM dNTP, 20 U RNasin, 2.5 µM random hexamers, and 50 U MMLV (Moloney murine leukemia virus) reverse transcriptase (RT). Reverse transcription was  
15 performed at room temperature for 10 min, and at 42°C for 60 min in a PE-480 thermal cycler, followed by further incubation at 95°C for 5 min to inactivate the RT. PCR was performed using 2 µl of the cDNA reaction in a final PCR reaction volume of 50 µl containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 2 mM MgCl<sub>2</sub>, 200 µM dNTP, 0.5 µM of each sense and antisense primer (SEQUENCE ID NO 12 and  
20 SEQUENCE ID NO 13, respectively), and 2.5 U of Taq polymerase. The reaction was incubated in an MJ Research Model PTC-200 as follows: 40 cycles of amplification (94°C, 20 sec; 58°C, 30 sec; 72°C, 30 sec); a final extension (72°C, 10 min); and a soak at 4°C.

C. PCR Fragment Analysis. The correct products were verified by size  
25 determination using gel electrophoresis with a SYBR® Green nucleic acid gel stain (Molecular Probes, Eugene, OR). Gels were stained with SYBR Green I at a 1:10,000 dilution in 1X TBE for 45 min. Gels were then de-stained in 1X TBE for 30 min. and imaged using a STORM imaging system (see Figures 4A and 4B). Figure 4A shows a DNA band at 201 bases from RNA of both normal (lanes 1-5)  
30 and cancerous (lanes 6-10) breast tissues. This band is indicative of a BS106 mRNA-specific RT-PCR product. As shown in Figure 4B, the product was not detected using RNA isolated from tissues of lung (lanes 1-5) or colon (lanes 6-10). Detection of a product comprising a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE  
35 ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof, indicated

the presence of BS106 mRNA, suggesting a diagnosis of a breast tissue disease or condition, such as breast cancer.

Example 9: OH-PCR

5           A. Probe selection and Labeling. Target-specific primers and probes are designed to detect the above-described target sequences by oligonucleotide hybridization PCR. International Publication Nos WO 92/10505, published 25 June 1992, and WO 92/11388, published 9 July 1992, teach methods for labeling oligonucleotides at their 5' and 3' ends, respectively. According to one known  
10       method for labeling an oligonucleotide, a label-phosphoramidite reagent is prepared and used to add the label to the oligonucleotide during its synthesis. For example, see N. T. Thuong et al., Tet. Letters 29(46):5905-5908 (1988); or J. S. Cohen et al., published U.S. Patent Application 07/246,688 (NTIS ORDER No. PAT-APPL-7-246,688) (1989). Preferably, probes are labeled at their 3' end to prevent  
15       participation in PCR and the formation of undesired extension products. For one step OH-PCR, the probe should have a  $T_m$  at least 15°C below the  $T_m$  of the primers. The primers and probes are utilized as specific binding members, with or without detectable labels, using standard phosphoramidite chemistry and/or post-synthetic labeling methods which are well-known to one skilled in the art.

20           B. One Step Oligo Hybridization PCR. OH-PCR is performed on a 200  $\mu$ l reaction containing 50 mM (N,N-bis[2-Hydroxyethyl]glycine), pH 8.15, 81.7 mM KOAc, 33.33 mM KOH, 0.01 mg/ml bovine serum albumin, 0.1 mM ethylene diaminetetraacetic acid, 0.02 mg/ml  $\text{NaN}_3$ , 8% w/v glycerol, 150  $\mu$ M each of dNTP, 0.25  $\mu$ M each primer, 3.75 nM probe, 5U rTth polymerase, 3.25 mM  $\text{Mn}(\text{OAc})_2$   
25       and 5  $\mu$ l blood equivalents of target (see Example 3). Since RNA and the rTth polymerase enzyme are unstable in the presence of  $\text{Mn}(\text{OAc})_2$ , the  $\text{Mn}(\text{OAc})_2$  should be added just before target addition. The reaction is incubated in a Perkin-Elmer Thermal Cycler 480. Optimal conditions for cDNA synthesis and thermal cycling can be readily determined by those skilled in the art. Conditions which may be  
30       found useful include cDNA synthesis (60°C, 30 min), 30-45 amplification cycles (94°C, 40 sec; 55-70°C, 60 sec), oligo-hybridization (97°C, 5 min; 15°C, 5 min; 15°C soak). The correct reaction product contains at least one of the strands of the PCR product and an internally hybridized probe.

C. OH-PCR Product Analysis. Amplified reaction products are detected on  
35       an LCx<sup>®</sup> analyzer system (available from Abbott Laboratories, Abbott Park, IL).

Briefly, the correct reaction product is captured by an antibody labeled microparticle at a capturable site on either the PCR product strand or the hybridization probe, and the complex is detected by binding of a detectable antibody conjugate to either a detectable site on the probe or the PCR strand. Only a complex containing a PCR strand hybridized with the internal probe is detectable. The detection of this complex then is indicative of the presence of BS106 mRNA, suggesting a diagnosis of a breast disease or condition, such as breast cancer.

Many other detection formats exist which can be used and/or modified by those skilled in the art to detect the presence of amplified or non-amplified BS106-derived nucleic acid sequences including, but not limited to, ligase chain reaction (LCR, Abbott Laboratories, Abbott Park, IL); Q-beta replicase (Gene-Trak™, Naperville, Illinois), branched chain reaction (Chiron, Emeryville, CA) and strand displacement assays (Becton Dickinson, Research Triangle Park, NC).

Example 10: Synthetic Peptide Production

Synthetic peptides represented by SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, respectively, were prepared based upon the predicted amino acid sequence of the open reading frame of BS106 (SEQUENCE ID NO 16) (see Example 1). All peptides were synthesized on a Symphony Peptide Synthesizer (available from Rainin Instrument Co, Emeryville California), using Fmoc chemistry, standard cycles and in-situ HBTU activation. Cleavage and deprotection conditions were as follows: a volume of 2.5 ml of cleavage reagent (77.5% v/v trifluoroacetic acid, 15% v/v ethanedithiol, 2.5% v/v water, 5% v/v thioanisole, 1-2% w/v phenol) was added to the resin, and agitated at room temperature for 2-4 hours. The filtrate was then removed and the peptide was precipitated from the cleavage reagent with cold diethyl ether. Each peptide was then filtered, purified via reverse-phase preparative HPLC using a water/acetonitrile/0.1% TFA gradient, and lyophilized. The product was confirmed by mass spectrometry (data not shown).

The purified peptides were mixed with adjuvant, and injected into rabbits (see Example 14). Alternatively, the purified peptides may be conjugated to Keyhole Limpet Hemocyanin with glutaraldehyde, mixed with adjuvant, and injected into rabbits (see Example 14).

Example 11a: Expression of Protein in a Cell Line Using Plasmid 577

A. Construction of a BS106 Expression Plasmid. Plasmid 577, described in U.S. patent application Serial No. 08/478,073, filed June 7, 1995 and incorporated herein by reference, has been constructed for the expression of secreted antigens in a permanent cell line. This plasmid contains the following DNA segments: (a) a 2.3 Kb fragment of pBR322 containing bacterial beta-lactamase and origin of DNA replication; (b) a 1.8 Kb cassette directing expression of a neomycin resistance gene under control of HSV-1 thymidine kinase promoter and poly-A addition signals; (c) a 1.9 Kb cassette directing expression of a dihydrofolate reductase gene under the control of an SV-40 promoter and poly-A addition signals; (d) a 3.5 Kb cassette directing expression of a rabbit immunoglobulin heavy chain signal sequence fused to a modified hepatitis C virus (HCV) E2 protein under the control of the Simian Virus 40 T-Ag promoter and transcription enhancer, the hepatitis B virus surface antigen (HBsAg) enhancer I followed by a fragment of Herpes Simplex Virus-1 (HSV-1) genome providing poly-A addition signals; and (e) a residual 0.7 Kb fragment of Simian Virus 40 genome late region of no function in this plasmid. All of the segments of the vector were assembled by standard methods known to those skilled in the art of molecular biology.

Plasmids for the expression of secretable BS106 proteins are constructed by replacing the hepatitis C virus E2 protein coding sequence in plasmid 577 with that of a BS106 polynucleotide sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof, as follows. Digestion of plasmid 577 with XbaI releases the hepatitis C virus E2 gene fragment. The resulting plasmid backbone allows insertion of the BS106 cDNA insert downstream of the rabbit immunoglobulin heavy chain signal sequence which directs the expressed proteins into the secretory pathway of the cell. The BS106 cDNA fragment is generated by PCR using standard procedures. Encoded in the sense PCR primer sequence is an XbaI site, immediately followed by a 12 nucleotide sequence that encodes the amino acid sequence Ser-Asn-Glu-Leu ("SNEL") to promote signal protease processing, efficient secretion and final product stability in culture fluids. Immediately following this 12 nucleotide sequence the primer contains nucleotides complementary to template sequences encoding amino acids of the BS106 gene. The antisense primer incorporates a sequence encoding the following eight amino acids just before the stop codons: Asp-Tyr-Lys-Asp-Asp-

Asp-Asp-Lys (SEQUENCE ID NO 21). Within this sequence is incorporated a recognition site to aid in analysis and purification of the BS106 protein product. A recognition site (termed "FLAG") that is recognized by a commercially available monoclonal antibody designated anti-FLAG M2 (Eastman Kodak, Co., New Haven, CT) can be utilized, as well as other comparable sequences and their corresponding antibodies. For example, PCR is performed using GeneAmp® reagents obtained from Perkin-Elmer-Cetus, as directed by the supplier's instructions. PCR primers are used at a final concentration of 0.5 µM. PCR is performed on the BS106 plasmid template in a 100 µl reaction for 35 cycles (94°C, 30 seconds; 55°C, 30 seconds; 72°C, 90 seconds) followed by an extension cycle of 72°C for 10 min.

B. Transfection of Dihydrofolate Reductase Deficient Chinese Hamster Ovary Cells. The plasmid described supra is transfected into CHO/dhfr- cells (DXB-111, Uriacio, et al., PNAS 77:4451-4466 (1980)). These cells are available from the A.T.C.C., 12301 Parklawn Drive, Rockville, MD 20852, under Accession No. CRL 9096. Transfection is carried out using the cationic liposome-mediated procedure described by P. L. Felgner et al., PNAS 84:7413-7417 (1987). Particularly, CHO/dhfr- cells are cultured in Ham's F-12 media supplemented with 10% fetal calf serum, L-glutamine (1 mM) and freshly seeded into a flask at a density of 5 - 8 x 10<sup>5</sup> cells per flask. The cells are grown to a confluency of between 60 and 80% for transfection. Twenty micrograms (20µg) of plasmid DNA is added to 1.5 ml of Opti-MEM I medium and 100 µl of Lipofectin Reagent (Gibco-BRL; Grand Island, NY) are added to a second 1.5 ml portion of Opti-MEM I media. The two solutions are mixed and incubated at room temperature for 20 min. After the culture medium is removed from the cells, the cells are rinsed 3 times with 5 ml of Opti-MEM I medium. The Opti-MEM I-Lipofection-plasmid DNA solution then is overlaid onto the cells. The cells are incubated for 3 h at 37°C, after which time the Opti-MEM I-Lipofectin-DNA solution is replaced with culture medium for an additional 24 h prior to selection.

C. Selection and Amplification. One day after transfection, cells are passaged 1:3 and incubated with dhfr/G418 selection medium (hereafter, "F-12 minus medium G"). Selection medium is Ham's F-12 with L-glutamine and without hypoxanthine, thymidine and glycine (JRH Biosciences, Lenexa, Kansas) and 300 µg per ml G418 (Gibco-BRL; Grand Island, NY). Media volume-to-surface area ratios of 5 ml per 25 cm<sup>2</sup> are maintained. After approximately two weeks,

451201-16029680



DHFR/G418 cells are expanded to allow passage and continuous maintenance in F-12 minus medium G.

Amplification of each of the transfected BS106 cDNA sequences is achieved by stepwise selection of DHFR<sup>+</sup>, G418<sup>+</sup> cells with methotrexate (reviewed by R. Schimke, Cell 37:705-713 (1984)). Cells are incubated with F-12 minus medium G containing 150 nM methotrexate (MTX) (Sigma, St. Louis, MO) for approximately two weeks until resistant colonies appear. Further gene amplification is achieved by selection of 150 nM adapted cells with 5  $\mu$ M MTX.

D. Antigen Production. F-12 minus medium G supplemented with 5  $\mu$ M MTX is overlaid onto just confluent monolayers for 12 to 24 h at 37°C in 5% CO<sub>2</sub>. The growth medium is removed and the cells are rinsed 3 times with Dulbecco's phosphate buffered saline (PBS) (with calcium and magnesium) (Gibco-BRL; Grand Island, NY) to remove the remaining media/serum which may be present. Cells then are incubated with VAS custom medium (VAS custom formulation with L-glutamine with HEPES without phenol red, available from JRH Bioscience; Lenexa, KS, product number 52-08678P), for 1 h at 37°C in 5% CO<sub>2</sub>. Cells then are overlaid with VAS for production at 5 ml per T flask. Medium is removed after seven days of incubation, retained, and then frozen to await purification with harvests 2, 3 and 4. The monolayers are overlaid with VAS for 3 more seven day harvests.

E. Analysis of Breast Tissue Gene BS106 Antigen Expression. Aliquots of VAS supernatants from the cells expressing the BS106 protein construct are analyzed, either by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) using standard methods and reagents known in the art (Laemmli discontinuous gels), or by mass spectrometry.

F. Purification. Purification of the BS106 protein containing the FLAG sequence is performed by immunoaffinity chromatography using an affinity matrix comprising anti-FLAG M2 monoclonal antibody covalently attached to agarose by hydrazide linkage (Eastman Kodak Co., New Haven, CT). Prior to affinity purification, protein in pooled VAS medium harvests from roller bottles is exchanged into 50 mM Tris-HCl (pH 7.5), 150 mM NaCl buffer using a Sephadex G-25 (Pharmacia Biotech Inc., Uppsala, Sweden) column. Protein in this buffer is applied to the anti-FLAG M2 antibody affinity column. Non-binding protein is eluted by washing the column with 50 mM Tris-HCl (pH 7.5), 150 mM NaCl buffer. Bound protein is eluted using an excess of FLAG peptide in 50 mM Tris-

HCl (pH 7.5), 150 mM NaCl. The excess FLAG peptide can be removed from the purified BS106 protein by gel electrophoresis or HPLC.

Although plasmid 577 is utilized in this example, it is known to those skilled in the art that other comparable expression systems, such as CMV, can be utilized  
5 herein with appropriate modifications in reagent and/or techniques and are within the skill of the ordinary artisan.

The largest cloned insert containing the coding region of the BS106 gene is sub-cloned into either (i) a eukaryotic expression vector which may contain, for example, a cytomegalovirus (CMV) promoter and/or protein fusible sequences  
10 which aid in protein expression and detection, or (ii) a bacterial expression vector containing a superoxide-dismutase (SOD) and CMP-KDO synthetase (CKS) or other protein fusion gene for expression of the protein sequence. Methods and vectors which are useful for the production of polypeptides which contain fusion sequences of SOD are described in EPO 0196056, published October 1, 1986, which is  
15 incorporated herein by reference and those containing fusion sequences of CKS are described in EPO Publication No. 0331961, published September 13, 1989, which publication is also incorporated herein by reference. This so-purified protein can be used in a variety of techniques, including but not limited to animal immunization studies, solid phase immunoassays, etc.

Example 11b: Expression of Protein in a Cell Line Using pcDNA3.1/Myc-His

A. Construction of a BS106 Expression Plasmid. Plasmid pcDNA3.1/Myc-His (Cat.# V855-20, Invitrogen, Carlsbad, CA) has been constructed, in the past, for the expression of secreted antigens by most mammalian cell lines. Expressed  
25 protein inserts are fused to a myc-his peptide tag. The myc-his tag (SEQUENCE ID NO 22) comprises a c-myc oncoprotein epitope and a polyhistidine sequence which are useful for the purification of an expressed fusion protein by using either anti-myc or anti-his affinity columns, or metalloprotein binding columns.

A plasmid for the expression of secretable BS106 protein was constructed by  
30 inserting a BS106 polynucleotide sequence from clone 1662885 into the pcDNA3.1/Myc-His vector. Prior to construction of the BS106 expression plasmid, the BS106 cDNA sequence was first cloned into a pCR®-Blunt vector. The BS106 cDNA fragment was generated by PCR performed using Stratagene® reagents obtained from Stratagene, as directed by the supplier's instructions. PCR primers  
35 are used at a final concentration of 0.5 µM. PCR using 5 U of pfu polymerase

(Stratagene, La Jolla, CA) was performed on the BS106 plasmid template (see Example 2) in a 50 µl reaction for 30 cycles (94°C, 1 min; 65°C, 1.5 min; 72°C, 3 min) followed by an extension cycle of 72°C for 8 min. The sense PCR primer sequence (SEQUENCE ID NO 14), comprises nucleotides which are identical to the pINCY vector directly upstream of the BS106 gene insert. The antisense primer (SEQUENCE ID NO 15), incorporates a 5' NotI restriction sequence and a sequence complementary to the 3' end of the BS106 cDNA insert just upstream of the 3'-most, in-frame stop codon. Five microliters (5 µl) of the resulting blunted-ended PCR product were ligated with 25 ng of linearized pCR®-Blunt vector (Invitrogen, Carlsbad, CA) interrupting the lethal ccdB gene of the vector. The resulting ligated vector was transformed into TOP10 *E. coli* (Invitrogen, Carlsbad, CA) using a One Shot™ transformation kit (Invitrogen, Carlsbad, CA) following supplier's directions. The transformed cells were grown on LB-Kan (50 µg/ml kanamycin) selection plates at 37°C. Only cells containing a plasmid with an interrupted ccdB gene grew after transformation (Grant, S.G.N., PNAS 87:4645-4649 (1990)). Transformed colonies were picked and grown in 3 ml of LB-Kan broth at 37°C. Plasmid DNA was isolated by using a QIAprep® (Qiagen Inc., Santa Clarita, CA) procedure, as directed by the supplier's instructions. The DNA was digested with EcoRI and NotI restriction enzymes to release the BS106 insert fragment. The fragment was electrophoresed on 1% Seakem® LE agarose (FMC, Rockland, ME)/0.5 µg/ml ethidium bromide/TE gel, visualized by UV illumination, excised and purified using QIAquick™ (Qiagen Inc., Santa Clarita, CA) procedures, as directed by the supplier's instructions.

The pcDNA3.1/Myc-His plasmid DNA was linearized by digestion with EcoRI and NotI, sites present in the polylinker region of the plasmid DNA. The BS106 purified fragment, supra, was ligated with the resulting plasmid DNA downstream from a CMV promoter, and transformed into DH5 alpha™ cells (GibcoBRL Gaithersburg, Md), as directed by the supplier's instructions. Briefly, 10 ng of pcDNA3.1/Myc-His containing the BS106 insert were added to 50 µl of competent DH5 alpha cells, and the contents were mixed gently. The mixture was incubated on ice for 30 min, heated for 20 sec at 37°C, and placed on ice for an additional 2 min. Upon addition of 0.95 ml of LB medium, the mixture was incubated for 1 h at 37°C while shaking at 225 rpm. The transformed cells then were plated onto 100 mm LB/Amp (50µg/ml ampicillin) plates and grown at 37°C. Colonies were picked and grown in 3 ml of LB/ampicillin broth. Plasmid DNA was

purified using a QIAprep kit. The presence of the insert was confirmed using restriction enzyme digestion and gel analysis (J. Sambrook et al., supra).

B. Transfection of Human Embryonic Kidney Cell 293 Cells. The BS106 expression plasmid described in Section A, supra, was retransformed into DH5 alpha cells, plated onto LB/ampicillin agar, and grown in 10 ml of LB/ampicillin broth, as described hereinabove. The plasmid was purified using a QIAfilter™ Maxi kit (Qiagen, Chatsworth, CA) and transfected into HEK293 cells (F.L. Graham et al., J. Gen. Vir. 36:59-72 (1977)). These cells are available from the A.T.C.C., 12301 Parklawn Drive, Rockville, MD 20852, under Accession No. CRL 1573.

Transfection was carried out using the cationic lipofectamine-mediated procedure described by P. Hawley-Nelson et al., Focus 15.73 (1993). Particularly, HEK293 cells were cultured in 10 ml DMEM media, supplemented with 10% fetal bovine serum (FBS), L-glutamine (2 mM), and freshly seeded into 12 x 100 mm culture plates at a density of  $8 \times 10^6$  cells per plate. The cells were grown at 37°C to a confluency of between 70% and 80% for transfection. Eight micrograms (8 µg) of plasmid DNA were added to 800 µl of Opti-MEM I® medium (Gibco-BRL, Grand Island, NY), and 48-96 µl of Lipofectamine™ Reagent (Gibco-BRL, Grand Island, NY) were added to a second 800 µl portion of Opti-MEM I medium. The two solutions were mixed and incubated at room temperature for 15-30 min. After the culture medium was removed from the cells, the cells were washed once with 10 ml of serum-free DMEM. The Opti-MEM I-Lipofectamine-plasmid DNA solution was diluted with 6.4 ml of serum-free DMEM and then overlaid onto the cells. The cells were incubated for 5 h at 37°C, after which time an additional 8 ml of DMEM with 20% FBS were added. After 18-24 h, the old medium was aspirated, and the cells were overlaid with 5 ml of fresh DMEM with 5% FBS. Supernatants and cell extracts were analyzed for BS106 gene activity 72 h after transfection.

C. Analysis of Breast Tissue Gene BS106 Antigen Expression. The culture supernatant, supra, was transferred to cryotubes and stored on ice. HEK293 cells were harvested by washing twice with 10 ml of cold Dulbecco's PBS and lysed by addition of 1.5 ml of CAT lysis buffer (Boehringer Mannheim, Indianapolis, IN), followed by incubation for 30 min at room temperature. Lysate was transferred to 1.7 ml polypropylene microfuge tubes and centrifuged at 1000 x g for 10 min. The supernatant was transferred to new cryotubes and stored on ice. Aliquots of supernatants from the cells and the lysate of the cells expressing the BS106 protein construct were analyzed for the presence of BS106 recombinant protein. The

aliquots were run on SDS-polyacrylamide gel electrophoresis (SDS-PAGE) using standard methods and reagents known in the art. (J. Sambrook et al., supra). Specifically, samples were adjusted to the desired protein concentration with Tricine buffer (Novex, San Diego, CA), mixed with an equal volume of 2X Tricine sample buffer (Novex, San Diego, CA), and heated for 5 minutes at 100°C in a thermal  
5 cyclor. Samples were then applied to a Novex 10-20% precast Tricine gel (Novex, San Diego, CA) for electrophoresis. Following electrophoresis, samples were electrophoretically transferred from the gels to Novex nitrocellulose membranes in Novex Tris-Glycine transfer buffer. The BS106 protein band was visualized using  
10 western blotting with anti-myc epitope monoclonal antibodies (Invitrogen, Carlsbad, CA) at a dilution of 1 to 5,000, and coupled with the Western Lights Plus chemiluminescence detection reagents (Tropix, Bedford, MA). The cell lysate showed the following two intense bands which stained specifically with the anti-myc monoclonal antibody: a sharp band at about 20 kD; and a broader band from about  
15 120-200 kD. The supernatant gave a single intense broad band from 30-45 kD, indicating differences in post-translational modification between the intracellular and the secreted form of BS106. Alternatively, anti-BS106 polyclonal serum (see Example 14) may be used to detect the recombinant BS106 protein, or the expressed BS106 recombinant protein may be analyzed by mass spectrometry (see Example  
20 12).

D. Purification. Purification of the BS106 recombinant protein containing the myc-his sequence is performed using the Xpress® affinity chromatography system (Invitrogen, Carlsbad, CA) containing a nickel-charged agarose resin which specifically binds polyhistidine residues. Supernatants from 10 x 100 mm plates,  
25 prepared as described supra, are pooled and passed over the nickel-charged column. Non-binding protein is eluted by washing the column with 50 mM Tris-HCl (pH 7.5)/150 mM NaCl buffer, leaving only the myc-his fusion proteins. Bound BS106 recombinant protein then is eluted from the column using either an excess of imidazole or histidine, or a low pH buffer. Alternatively, the recombinant protein  
30 can also be purified by binding at the myc-his sequence to an affinity column consisting of either anti-myc or anti-histidine monoclonal antibodies conjugated through a hydrazide or other linkage to an agarose resin and eluting with an excess of myc peptide or histidine, respectively.

The purified recombinant protein can then be covalently cross-linked to a  
35 solid phase, such as N-hydroxysuccinimide-activated sepharose columns (Pharmacia

467001-1609500

Biotech, Piscataway, NJ), as directed by supplier's instructions. These columns containing covalently linked BS106 recombinant protein, can then be used to purify anti-BS106 antibodies from rabbit or mouse sera (see Examples 13 and 14).

E. Coating Microtiter Plates with BS106 Expressed Proteins. Supernatant  
5 from a 100 mm plate, as described supra, is diluted in an appropriate volume of PBS. Then, 100 µl of the resulting mixture is placed into each well of a Reacti-Bind™ metal chelate microtiter plate (Pierce, Rockford, IL), incubated at room temperature while shaking, and followed by three washes with 200 µl each of PBS with 0.05% Tween® 20. The prepared microtiter plate can then be used to screen  
10 polyclonal antisera for the presence of BS106 antibodies (see Example 17).

Although pcDNA3.1/Myc-His is utilized in this example, it is known to those skilled in the art that other comparable expression systems can be utilized herein with appropriate modifications in reagent and/or techniques and are within the skill of one of ordinary skill in the art. The largest cloned insert containing the  
15 coding region of the BS106 gene is sub-cloned into either (i) a eukaryotic expression vector which may contain, for example, a cytomegalovirus (CMV) promoter and/or protein fusible sequences which aid in protein expression and detection, or (ii) a bacterial expression vector containing a superoxide-dismutase (SOD) and CMP-KDO synthetase (CKS) or other protein fusion gene for expression of the protein  
20 sequence. Methods and vectors which are useful for the production of polypeptides which contain fusion sequences of SOD are described in published EPO application No. EP 0 196 056, published October 1, 1986, which is incorporated herein by reference, and vectors containing fusion sequences of CKS are described in  
25 published EPO application No. EP 0 331 961, published September 13, 1989, which publication is also incorporated herein by reference. The purified protein can be used in a variety of techniques, including, but not limited to, animal immunization studies, solid phase immunoassays, and the like.

Example 12: Chemical Analysis of Breast Tissue Proteins

30 A. Analysis of Tryptic Peptide Fragments Using MS. Sera from patients with breast disease such as breast cancer, sera from patients with no breast disease, extracts of breast tissues or cells from patients with breast disease such as breast cancer, extracts of breast tissues or cells from patients with no breast disease, and extracts of tissues or cells from other non-diseased or diseased organs of patients,  
35 are run on a polyacrylamide gel using standard procedures and stained with

Coomassie Blue. Sections of the gel suspected of containing the unknown polypeptide are excised and subjected to an in-gel reduction, acetamidation and tryptic digestion. P. Jenö et al, Anal. Bio. 224:451-455 (1995) and J. Rosenfeld et al, Anal. Bio. 203:173-179 (1992). The gel sections are washed with 100 mM

5  $\text{NH}_4\text{HCO}_3$  and acetonitrile. The shrunken gel pieces are swollen in digestion buffer (50 mM  $\text{NH}_4\text{HCO}_3$ , 5 mM  $\text{CaCl}_2$  and 12.5  $\mu\text{g/ml}$  trypsin) at 4°C for 45 min. The supernatant is aspirated and replaced with 5 to 10  $\mu\text{l}$  of digestion buffer without trypsin and allowed to incubate overnight at 37°C. Peptides are extracted with 3 changes of 5% formic acid and acetonitrile and evaporated to dryness. The peptides

10 are adsorbed to approximately 0.1  $\mu\text{l}$  of POROS R2 sorbent (Perseptive Biosystems, Framingham, Massachusetts) trapped in the tip of a drawn gas chromatography capillary tube by dissolving them in 10  $\mu\text{l}$  of 5% formic acid and passing it through the capillary. The adsorbed peptides are washed with water and eluted with 5% formic acid in 60% methanol. The eluant is passed directly into the spraying

15 capillary of an API III mass spectrometer (Perkin-Elmer Sciex, Thornhill, Ontario, Canada) for analysis by nano-electrospray mass spectrometry. M. Wilm et al., Int. J. Mass Spectrom. Ion Process 136:167-180 (1994) and M. Wilm et al., Anal. Chem. 66:1-8 (1994). The masses of the tryptic peptides are determined from the mass spectrum obtained off the first quadrupole. Masses corresponding to predicted

20 peptides can be further analyzed in MS/MS mode to give the amino acid sequence of the peptide.

B. Peptide Fragment Analysis Using LC/MS. The presence of polypeptides predicted from mRNA sequences found in hyperplastic disease tissues also can be confirmed using liquid chromatography/tandem mass spectrometry (LC/MS/MS).

25 D. Hess et al., METHODS, A Companion to Methods in Enzymology 6:227-238 (1994). The serum specimen or tumor extract from the patient is denatured with SDS and reduced with dithiothreitol (1.5 mg/ml) for 30 min at 90°C followed by alkylation with iodoacetamide (4 mg/ml) for 15 min at 25°C. Following acrylamide electrophoresis, the polypeptides are electroblotted to a cationic membrane and

30 stained with Coomassie Blue. Following staining, the membranes are washed and sections thought to contain the unknown polypeptides are cut out and dissected into small pieces. The membranes are placed in 500  $\mu\text{l}$  microcentrifuge tubes and immersed in 10 to 20  $\mu\text{l}$  of proteolytic digestion buffer (100 mM Tris-HCl, pH 8.2, containing 0.1 M NaCl, 10% acetonitrile, 2 mM  $\text{CaCl}_2$  and 5  $\mu\text{g/ml}$  trypsin) (Sigma,

35 St. Louis, MO). After 15 h at 37°C, 3  $\mu\text{l}$  of saturated urea and 1  $\mu\text{l}$  of 100  $\mu\text{g/ml}$

trypsin are added and incubated for an additional 5 h at 37°C. The digestion mixture is acidified with 3 µl of 10% trifluoroacetic acid and centrifuged to separate supernatant from membrane. The supernatant is injected directly onto a microbore, reverse phase HPLC column and eluted with a linear gradient of acetonitrile in  
5 0.05% trifluoroacetic acid. The eluate is fed directly into an electrospray mass spectrometer, after passing through a stream splitter if necessary to adjust the volume of material. The data is analyzed following the procedures set forth in Example 12, Section A.

10 Example 13: Gene Immunization Protocol

A. In Vivo Antigen Expression. Gene immunization circumvents protein purification steps by directly expressing an antigen in vivo after inoculation of the appropriate expression vector. Also, production of antigen by this method may allow correct protein folding and glycosylation since the protein is produced in  
15 mammalian tissue. The method utilizes insertion of the gene sequence into a plasmid which contains a CMV promoter, expansion and purification of the plasmid and injection of the plasmid DNA into the muscle tissue of an animal. Preferred animals include mice and rabbits. See, for example, H. Davis et al., Human Molecular Genetics 2:1847-1851 (1993). After one or two booster immunizations, the animal  
20 can then be bled, ascites fluid collected, or the animal's spleen can be harvested for production of hybridomas.

B. Plasmid Preparation and Purification. BS106 cDNA insert was released from the BS106 vector described in Example 11b by digestion with EcoRI and NotI restriction enzymes. The digested plasmid fragments were electrophoresed on a 1%  
25 Seakem KE agarose/0.5 µg/ml ethidium bromide/TE gel and the bands were visualized by UV illumination. The insert fragment was excised from the gel and purified using the QIAquick procedure, described supra. The fragment was ligated into an EcoRI/NotI digested pcDNA3.1 vector (Invitrogen, Carlsbad, CA) and transformed into DH5 alpha cells as described supra. The plasmid DNA was  
30 purified from the bacterial lysate using a QIAprep column. All these techniques are familiar to one of ordinary skill in the art of molecular biology.

C. Immunization Protocol. Anesthetized animals are immunized intramuscularly with 0.1-100 µg of the purified plasmid diluted in PBS or other DNA uptake enhancers (Cardiotoxin, 25% sucrose). See, for example, H. Davis et

46299680



al, Human Gene Therapy 4:733-740 (1993); and P. W. Wolff et al, Biotechniques 11:474-485 (1991). One to two booster injections are given at monthly intervals.

D. Testing and Use of Antiserum. Animals are bled and the resultant sera tested for antibody using peptides synthesized from the known gene sequence (see Example 16) using techniques known in the art, such as western blotting or EIA techniques. Antisera produced by this method can then be used to detect the presence of the antigen in a patient's tissue or cell extract, or in a patient's serum, by ELISA or Western blotting techniques, such as those described in Examples 15 through 18.

#### Example 14: Production of Antibodies Against BS106

A. Production of Polyclonal Antisera. Antiserum against BS106 is prepared by injecting rabbits with peptides whose sequences are derived from that of the predicted amino acid sequence of the BS106 consensus sequence (SEQUENCE ID NO 4). The synthesis of peptides (SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, and SEQUENCE ID NO 20) is described in Example 10.

1. Peptide Conjugation. Peptide is conjugated to maleimide activated keyhole limpet hemocyanin (KLH, commercially available as Imject<sup>®</sup>, available from Pierce Chemical Company, Rockford, IL). Imject<sup>®</sup> contains about 250 moles of reactive maleimide groups per mole of hemocyanin. The activated KLH is dissolved in phosphate buffered saline (PBS, pH 8.4) at a concentration of about 7.7 mg/ml. The peptide is conjugated through cysteines occurring in the peptide sequence, or to a cysteine previously added to the synthesized peptide in order to provide a point of attachment. The peptide is dissolved in dimethyl sulfoxide (DMSO, Sigma Chemical Company, St. Louis, MO) and reacted with the activated KLH at a mole ratio of about 1.5 moles of peptide per mole of reactive maleimide attached to the KLH. A procedure for the conjugation of peptide is provided hereinbelow. It is known to the ordinary artisan that the amounts, times and conditions of such a procedure can be varied to optimize peptide conjugation.

The conjugation reaction described hereinbelow is based on obtaining 3 mg of KLH peptide conjugate ("conjugated peptide"), which contains about 0.77  $\mu$ moles of reactive maleimide groups. This quantity of peptide conjugate usually is adequate for one primary injection and four booster injections for production of polyclonal antisera in a rabbit. Briefly, peptide is dissolved in DMSO at a

concentration of 1.16  $\mu$ moles/100  $\mu$ l of DMSO. One hundred microliters (100  $\mu$ l) of the DMSO solution are added to 380  $\mu$ l of the activated KLH solution prepared as described hereinabove, and 20  $\mu$ l of PBS (pH 8.4) is added to bring the volume to 500  $\mu$ l. The reaction is incubated overnight at room temperature with stirring. The  
5 extent of reaction is determined by measuring the amount of unreacted thiol in the reaction mixture. The difference between the starting concentration of thiol and the final concentration is assumed to be the concentration of peptide which has coupled to the activated KLH. The amount of remaining thiol is measured using Ellman's reagent (5,5'-dithiobis(2-nitrobenzoic acid), Pierce Chemical Company, Rockford,  
10 IL). Cysteine standards are made at a concentration of 0, 0.1, 0.5, 2, 5 and 20 mM by dissolving 35 mg of cysteine HCl (Pierce Chemical Company, Rockford, IL) in 10 ml of PBS (pH 7.2) and diluting the stock solution to the desired concentration. The photometric determination of the concentration of thiol is accomplished by  
15 placing 200  $\mu$ l of PBS (pH 8.4) in each well of an Immulon 2<sup>®</sup> microwell plate (Dynex Technologies, Chantilly, VA). Next, 10  $\mu$ l of standard or reaction mixture is added to each well. Finally, 20  $\mu$ l of Ellman's reagent at a concentration of 1 mg/ml in PBS (pH 8.4) is added to each well. The wells are incubated for 10 minutes at room temperature, and the absorbance of all wells is read at 415 nm with  
20 a microplate reader (such as the BioRad Model 3550, BioRad, Richmond, CA). The absorbance of the standards is used to construct a standard curve and the thiol concentration of the reaction mixture is determined from the standard curve. A decrease in the concentration of free thiol is indicative of a successful conjugation reaction. In addition, calculation of free thiol in the peptide solution, prior to  
25 addition of the maleimide activated KLH and upon completion of the reaction, allows determination of the substitution ratio of moles of peptide/mole of KLH for each peptide x KLH conjugate. In all cases, the reaction would be taken to completion, and there would be approximately 250 peptides/KLH molecule for each peptide conjugate prepared. Any unreacted peptide is removed by dialysis against PBS (pH  
30 7.2) at room temperature for 6 hours. The conjugate is stored at 2-8°C if it is to be used immediately; otherwise, it is stored at -20°C or colder.

2. Animal Immunization. Female white New Zealand rabbits weighing 2 kg or more were used for raising polyclonal antiserum. One animal was immunized per peptide (SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, and SEQUENCE ID NO 20, respectively). One week prior

to the first immunization, 5 to 10 ml of blood were obtained from the animal to serve as a non-immune prebled sample.

Each of the peptides, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, and SEQUENCE ID NO 20, was used to prepare the primary immunogen by emulsifying 0.5 ml of the peptide at a concentration of 2 mg/ml in PBS (pH 7.2) with 0.5 ml of complete Freund's adjuvant (CFA) (Difco, Detroit, MI). The immunogen was injected into several sites of the animal via subcutaneous, intraperitoneal, and/or intramuscular routes of administration. Four weeks following the primary immunization, a booster immunization was administered. The immunogen used for the booster immunization dose was prepared by emulsifying 0.5 ml of the same peptide used for the primary immunogen, except that the peptide now was diluted to 1 mg/ml with 0.5 ml of incomplete Freund's adjuvant (IFA) (Difco, Detroit, MI). Again, the booster dose was administered into several sites and utilized subcutaneous, intraperitoneal and intramuscular types of injections. The animal was bled (5 ml) two weeks after the booster immunization and the serum was tested for immunoreactivity to the peptide, as described below. The booster and bleed schedule is repeated at 4 week intervals until an adequate titer is obtained. The titer or concentration of antiserum is determined by microtiter EIA as described in Example 17, below. In addition, apparent affinity values [ $K_d$ (app)] are determined for some of the antisera (see Example 17). An antibody titer of 1:500 or greater is considered an adequate titer for further use and study.

#### B. Production of Monoclonal Antibody.

1. Immunization Protocol. Mice are immunized using immunogens prepared as described hereinabove, except that the amount of the unconjugated or conjugated peptide for monoclonal antibody production in mice is one-tenth the amount used to produce polyclonal antisera in rabbits. Thus, the primary immunogen consists of 100  $\mu$ g of unconjugated or conjugated peptide in 0.1 ml of CFA emulsion; while the immunogen used for booster immunizations consists of 50  $\mu$ g of unconjugated or conjugated peptide in 0.1 ml of IFA. Hybridomas for the generation of monoclonal antibodies are prepared and screened using standard techniques. The methods used for monoclonal antibody development follow procedures known in the art such as those detailed in Kohler and Milstein, Nature 256:494 (1975) and reviewed in J.G.R. Hurrel, ed., Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press, Inc., Boca Raton, FL

(1982). Another method of monoclonal antibody development which is based on the Kohler and Milstein method is that of L.T. Mimms et al., Virology 176:604-619 (1990), which is incorporated herein by reference.

The immunization regimen (per mouse) consists of a primary immunization  
5 with additional booster immunizations. The primary immunogen used for the  
primary immunization consists of 100 µg of unconjugated or conjugated peptide in  
50 µl of PBS (pH 7.2) previously emulsified in 50 µl of CFA. Booster  
immunizations performed at approximately two weeks and four weeks post primary  
immunization consist of 50 µg of unconjugated or conjugated peptide in 50 µl of  
10 PBS (pH 7.2) emulsified with 50 µl IFA. A total of 100 µl of this immunogen is  
inoculated intraperitoneally and subcutaneously into each mouse. Individual mice  
are screened for immune response by microtiter plate enzyme immunoassay (EIA) as  
described in Example 17 approximately four weeks after the third immunization.  
Mice are inoculated either intravenously, intrasplenically or intraperitoneally with 50  
15 µg of unconjugated or conjugated peptide in PBS (pH 7.2) approximately fifteen  
weeks after the third immunization..

Three days after this intravenous boost, splenocytes are fused with, for  
example, Sp2/0-Ag14 myeloma cells (Milstein Laboratories, England) using the  
polyethylene glycol (PEG) method. The fusions are cultured in Iscove's Modified  
20 Dulbecco's Medium (IMDM) containing 10% fetal calf serum (FCS), plus 1%  
hypoxanthine, aminopterin and thymidine (HAT). Bulk cultures are screened by  
microtiter plate EIA following the protocol in Example 17. Clones reactive with the  
peptide used an immunogen and non-reactive with other peptides (i.e., peptides of  
BS106 not used as the immunogen) are selected for final expansion. Clones thus  
25 selected are expanded, aliquoted and frozen in IMDM containing 10% FCS and 10%  
dimethyl-sulfoxide.

## 2. Production of Ascites Fluid Containing Monoclonal Antibodies.

Frozen hybridoma cells prepared as described hereinabove are thawed and placed  
into expansion culture. Viable hybridoma cells are inoculated intraperitoneally into  
30 Pristane treated mice. Ascitic fluid is removed from the mice, pooled, filtered  
through a 0.2 µ filter and subjected to an immunoglobulin class G (IgG) analysis to  
determine the volume of the Protein A column required for the purification.

## 3. Purification of Monoclonal Antibodies From Ascites Fluid.

Briefly, filtered and thawed ascites fluid is mixed with an equal volume of Protein A  
35 sepharose binding buffer (1.5 M glycine, 3.0 M NaCl, pH 8.9) and refiltered

through a 0.2  $\mu$  filter. The volume of the Protein A column is determined by the quantity of IgG present in the ascites fluid. The eluate then is dialyzed against PBS (pH 7.2) overnight at 2-8°C. The dialyzed monoclonal antibody is sterile filtered and dispensed in aliquots. The immunoreactivity of the purified monoclonal antibody is confirmed by determining its ability to specifically bind to the peptide used as the immunogen by use of the EIA microtiter plate assay procedure of Example 17. The specificity of the purified monoclonal antibody is confirmed by determining its lack of binding to irrelevant peptides such as peptides of BS106 not used as the immunogen. The purified anti-BS106 monoclonal thus prepared and characterized is placed at either 2-8°C for short term storage or at -80°C for long term storage.

4. Further Characterization of Monoclonal Antibody. The isotype and subtype of the monoclonal antibody produced as described hereinabove can be determined using commercially available kits (available from Amersham, Inc., Arlington Heights, IL). Stability testing also can be performed on the monoclonal antibody by placing an aliquot of the monoclonal antibody in continuous storage at 2-8°C and assaying optical density (OD) readings throughout the course of a given period of time.

C. Use of Recombinant Proteins as Immunogens. It is within the scope of the present invention that recombinant proteins made as described herein can be utilized as immunogens in the production of polyclonal and monoclonal antibodies, with corresponding changes in reagents and techniques known to those skilled in the art.

Example 15: Purification of Serum Antibodies Which Specifically Bind to BS106 Peptides

Immune sera, obtained as described hereinabove in Examples 13 and/or 14, is affinity purified using immobilized synthetic peptides prepared as described in Example 10, or recombinant proteins prepared as described in Example 11. An IgG fraction of the antiserum is obtained by passing the diluted, crude antiserum over a Protein A column (Affi-Gel protein A, Bio-Rad, Hercules, CA). Elution with a buffer (Binding Buffer, supplied by the manufacturer) removes substantially all proteins that are not immunoglobulins. Elution with 0.1M buffered glycine (pH 3) gives an immunoglobulin preparation that is substantially free of albumin and other serum proteins.

Immunoaffinity chromatography is performed to obtain a preparation with a higher fraction of specific antigen-binding antibody. The peptide used to raise the antiserum is immobilized on a chromatography resin, and the specific antibodies directed against its epitopes are adsorbed to the resin. After washing away non-binding components, the specific antibodies are eluted with 0.1 M glycine buffer (pH 2.3). Antibody fractions are immediately neutralized with 1.0 M Tris buffer (pH 8.0) to preserve immunoreactivity. The chromatography resin chosen depends on the reactive groups present in the peptide. If the peptide has an amino group, a resin such as Affi-Gel 10 or Affi-Gel 15 is used (Bio-Rad, Hercules, CA). If coupling through a carboxy group on the peptide is desired, Affi-Gel 102 can be used (Bio-Rad, Hercules, CA). If the peptide has a free sulfhydryl group, an organomercurial resin such as Affi-Gel 501 (Bio-Rad, Hercules, CA) or SulkfoLink™ (Pierce, Rockford, IL) can be used. The amount of peptide immobilized on the resin can be determined using Nano Orange™ (Molecular Probes, Eugene, OR).

Alternatively, spleens can be harvested and used in the production of hybridomas to produce monoclonal antibodies following routine methods known in the art as described hereinabove.

#### 20                    Example 16: Western Blotting of Tissue Samples

Protein extracts are prepared by homogenizing tissue samples in 0.1M Tris-HCl (pH 7.5), 15% (w/v) glycerol, 0.2 mM EDTA, 1.0 mM 1,4-dithiothreitol, 10 µg/ml leupeptin and 1.0 mM phenylmethylsulfonylfluoride (Kain et al., Biotechniques, 17:982 (1994)). Following homogenization, the homogenates are centrifuged at 4°C for 5 minutes to separate supernate from debris. For protein quantitation, 3-10 µl of supernate are added to 1.5 ml of bicinchoninic acid reagent (Sigma, St. Louis, MO), and the resulting absorbance at 562 nm is measured.

For SDS-PAGE, samples are adjusted to desired protein concentration with Tricine Buffer (Novex, San Diego, CA), mixed with an equal volume of 2X Tricine sample buffer (Novex, San Diego, CA), and heated for 5 minutes at 100°C in a thermal cycler. Samples are then applied to a Novex 10-20% Precast Tricine Gel for electrophoresis. Following electrophoresis, samples are transferred from the gels to nitrocellulose membranes in Novex Tris-Glycine Transfer buffer. Membranes are then probed with specific anti-peptide antibodies using the reagents and procedures provided in the Western Lights or Western Lights Plus (Tropix, Bedford, MA)

chemiluminescence detection kits. Chemiluminescent bands are visualized by exposing the developed membranes to Hyperfilm ECL (Amersham, Arlington Heights, IL).

Competition experiments are carried out in an analogous manner as above, with the following exception; the primary antibodies (anti-peptide polyclonal antisera) are pre-incubated for 30 minutes at room temperature with varying concentrations of peptide immunogen prior to exposure to the nitrocellulose filter. Development of the Western is performed as above.

After visualization of the bands on film, the bands can also be visualized directly on the membranes by the addition and development of a chromogenic substrate such as 5-bromo-4-chloro-3-indolyl phosphate (BCIP). This chromogenic solution contains 0.016% BCIP in a solution containing 100 mM NaCl, 5 mM MgCl<sub>2</sub> and 100 mM Tris-HCl (pH 9.5). The filter is incubated in the solution at room temperature until the bands develop to the desired intensity. Molecular mass determination is made based upon the mobility of pre-stained molecular weight standards (Novex, San Diego, CA) or biotinylated molecular weight standards (Tropix, Bedford, MA).

#### Example 17: EIA Microtiter Plate Assay

The immunoreactivity of antiserum preferably obtained from rabbits or mice as described in Example 13 or Example 14 is determined by means of a microtiter plate EIA, as follows. Synthetic peptides prepared as described in Example 10 or recombinant proteins prepared as described in Example 11 are dissolved in 50 mM carbonate buffer (pH 9.6) to a final concentration of 2 µg/ml. Next, 100 µl of the peptide or protein solution is placed in each well of an Immulon 2<sup>®</sup> microtiter plate (Dynex Technologies, Chantilly, VA). The plate is incubated overnight at room temperature and then washed four times with deionized water. The wells are blocked by adding 125 µl of a suitable protein blocking agent, such as Superblock<sup>®</sup> (Pierce Chemical Company, Rockford, IL), in phosphate buffered saline (PBS, pH 7.4) to each well and then immediately discarding the solution. This blocking procedure is performed three times. Antiserum obtained from immunized rabbits or mice prepared as previously described is diluted in a protein blocking agent (e.g., a 3% Superblock<sup>®</sup> solution) in PBS containing 0.05% Tween-20<sup>®</sup> (monolaurate polyoxyethylene ether) (Sigma Chemical Company, St. Louis, MO) and 0.05% sodium azide at dilutions of 1:500, 1:2500, 1:12,500, 1:62,500 and 1:312,500 and

placed in each well of the coated microtiter plate. The wells then are incubated for three hours at room temperature. Each well is washed four times with deionized water. One hundred  $\mu$ l of alkaline phosphatase-conjugated goat anti-rabbit IgG or goat anti-mouse IgG antiserum (Southern Biotech, Birmingham, AB), diluted  
5 1:2000 in 3% Superblock<sup>®</sup> solution in phosphate buffered saline containing 0.05% Tween 20<sup>®</sup> and 0.05% sodium azide, is added to each well. The wells are incubated for two hours at room temperature. Next, each well is washed four times with deionized water. One hundred microliters (100  $\mu$ l) of paranitrophenyl phosphate substrate (Kirkegaard and Perry Laboratories, Gaithersburg, MD) then is added to  
10 each well. The wells are incubated for thirty minutes at room temperature. The absorbance at 405 nm is read of each well. Positive reactions are identified by an increase in absorbance at 405 nm in the test well above that absorbance given by a non-immune serum (negative control). A positive reaction is indicative of the presence of detectable anti-BS106 antibodies.

15 In addition to titers, apparent affinities [ $K_d$ (app)] may also be determined for some of the anti-peptide antisera. EIA microtiter plate assay results can be used to derive the apparent dissociation constants ( $K_d$ ) based on an analog of the Michaelis-Menten equation (V. Van Heyningen, Methods in Enzymology, Vol. 121, p. 472 (1986) and further described in X. Qiu, et al, Journal of Immunology, Vol. 156, p.  
20 3350 (1996)):

$$[Ag-Ab] = [Ag-Ab]_{max} \times \frac{[Ab]}{[Ab] + K_d}$$

25 Where [Ag-Ab] is the antigen-antibody complex concentration,  $[Ag-Ab]_{max}$  is the maximum complex concentration, [Ab] is the antibody concentration, and  $K_d$  is the dissociation constant. During the curve fitting, the [Ag-Ab] is replaced with the background subtracted value of the OD<sub>405nm</sub> at the given concentration of Ab. Both  $K_d$  and  $[OD_{405nm}]_{max}$ , which corresponds to the  $[Ag-Ab]_{max}$ , are treated as fitted  
30 parameters. The software program Origin can be used for the curve fitting.

#### Example 18: Coating of Solid Phase Particles

A. Coating of Microparticles with Antibodies Which Specifically Bind to BS106 Antigen. Affinity purified antibodies which specifically bind to BS106 protein (see Example 15) are coated onto microparticles of polystyrene, carboxylated  
35 polystyrene, polymethylacrylate or similar particles having a radius in the range of



about 0.1 to 20  $\mu\text{m}$ . Microparticles may be either passively or actively coated. One coating method comprises coating EDAC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (Aldrich Chemical Co., Milwaukee, WI) activated carboxylated latex microparticles with antibodies which specifically bind to BS106 protein, as follows. Briefly, a final 0.375% solid suspension of resin washed carboxylated latex microparticles (available from Bangs Laboratories, Carmel, IN or Serodyn, Indianapolis, IN) are mixed in a solution containing 50 mM MES buffer, pH 4.0 and 150 mg/l of affinity purified anti-BS106 antibody (see Example 14) for 15 min in an appropriate container. EDAC coupling agent is added to a final concentration of 5.5  $\mu\text{g}/\text{ml}$  to the mixture and mixed for 2.5 h at room temperature.

The microparticles then are washed with 8 volumes of a Tween 20®/sodium phosphate wash buffer (pH 7.2) by tangential flow filtration using a 0.2  $\mu\text{m}$  Microgon Filtration module. Washed microparticles are stored in an appropriate buffer which usually contains a dilute surfactant and irrelevant protein as a blocking agent, until needed.

B. Coating of 1/4 Inch Beads. Antibodies which specifically bind to BS106-antigen also may be coated on the surface of 1/4 inch polystyrene beads by routine methods known in the art (Snitman et al, US Patent 5,273,882, incorporated herein by reference) and used in competitive binding or EIA sandwich assays.

Polystyrene beads first are cleaned by ultrasonication for about 15 seconds in 10 mM  $\text{NaHCO}_3$  buffer at pH 8.0. The beads then are washed in deionized water until all fines are removed. Beads then are immersed in an antibody solution in 10 mM carbonate buffer, pH 8 to 9.5. The antibody solution can be as dilute as 1  $\mu\text{g}/\text{ml}$  in the case of high affinity monoclonal antibodies or as concentrated as about 500  $\mu\text{g}/\text{ml}$  for polyclonal antibodies which have not been affinity purified. Beads are coated for at least 12 hours at room temperature, and then they are washed with deionized water. Beads may be air dried or stored wet (in PBS, pH 7.4). They also may be overcoated with protein stabilizers (such as sucrose) or protein blocking agents used as non-specific binding blockers (such as irrelevant proteins, Carnation skim milk, Superblock®, or the like).

#### Example 19: Microparticle Enzyme Immunoassay (MEIA)

BS106 antigens are detected in patient test samples by performing a standard antigen competition EIA or antibody sandwich EIA and utilizing a solid phase such

as microparticles (MEIA). The assay can be performed on an automated analyzer such as the IMx<sup>®</sup> Analyzer (Abbott Laboratories, Abbott Park, IL).

A. Antibody Sandwich EIA. Briefly, samples suspected of containing BS106 antigen are incubated in the presence of anti-BS106 antibody-coated  
5 microparticles (prepared as described in Example 17) in order to form antigen/antibody complexes. The microparticles then are washed and an indicator reagent comprising an antibody conjugated to a signal generating compound (i.e., enzymes such as alkaline phosphatase or horseradish peroxide) is added to the antigen/antibody complexes or the microparticles and incubated. The microparticles  
10 are washed and the bound antibody/antigen/antibody complexes are detected by adding a substrate (e.g., 4-methyl umbelliferyl phosphate (MUP), or OPD/peroxide, respectively), that reacts with the signal generating compound to generate a measurable signal. An elevated signal in the test sample, compared to the signal generated by a negative control, detects the presence of BS106 antigen. The  
15 presence of BS106 antigen in the test sample is indicative of a diagnosis of a breast disease or condition, such as breast cancer.

B. Competitive Binding Assay. The competitive binding assay uses a peptide or protein that generates a measurable signal when the labeled peptide is contacted with an anti-peptide antibody coated microparticle. This assay can be  
20 performed on the IMx<sup>®</sup> Analyzer (available from Abbott Laboratories, Abbott Park, IL). The labeled peptide is added to the BS106 antibody-coated microparticles (prepared as described in Example 17) in the presence of a test sample suspected of containing BS106 antigen, and incubated for a time and under conditions sufficient to form labeled BS106 peptide (or labeled protein) / bound antibody complexes  
25 and/or patient BS106 antigen / bound antibody complexes. The BS106 antigen in the test sample competes with the labeled BS106 peptide (or BS106 protein) for binding sites on the microparticle. BS106 antigen in the test sample results in a lowered binding of labeled peptide and antibody coated microparticles in the assay since antigen in the test sample and the BS106 peptide or BS106 protein compete for  
30 antibody binding sites. A lowered signal (compared to a control) indicates the presence of BS106 antigen in the test sample. The presence of BS106 antigen suggests the diagnosis of a breast disease or condition, such as breast cancer.

The BS106 polynucleotides and the proteins encoded thereby which are provided and discussed hereinabove are useful as markers of breast tissue disease,  
35 especially breast cancer. Tests based upon the appearance of this marker in a test

sample such as blood, plasma or serum can provide low cost, non-invasive, diagnostic information to aid the physician to make a diagnosis of cancer, to help select a therapy protocol, or to monitor the success of a chosen therapy. This marker may appear in readily accessible body fluids such as blood, urine or stool as antigens derived from the diseased tissue which are detectable by immunological methods. 5 This marker may be elevated in a disease state, altered in a disease state, or be a normal protein of the breast which appears in an inappropriate body compartment.

4629580

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: Billing-Medel, Patricia A.  
Cohen, Maurice  
Colpitts, Tracey L.  
Friedman, Paula N.  
Gordon, Julian  
Granados, Edward N.  
Hodges, Steven C.  
Klass, Michael R.  
Kratochvil, Jon D.  
Roberts-Rapp, Lisa  
Russell, John C.  
Stroupe, Steven D.
- (ii) TITLE OF THE INVENTION: REAGENTS AND METHODS USEFUL  
FOR DETECTING DISEASES OF THE BREAST
- (iii) NUMBER OF SEQUENCES: 22
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Abbott Laboratories  
(B) STREET: 100 Abbott Park Road  
(C) CITY: Abbott Park  
(D) STATE: IL  
(E) COUNTRY: USA  
(F) ZIP: 60064-3500
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Diskette  
(B) COMPUTER: IBM Compatible  
(C) OPERATING SYSTEM: DOS  
(D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: 08/742,067  
(B) FILING DATE: 31-OCT-1996
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Becker, Cheryl L  
(B) REGISTRATION NUMBER: 35,441  
(C) REFERENCE/DOCKET NUMBER: 5995.US.P1
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: 847/935-1729  
(B) TELEFAX: 847/938-2623  
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 229 base pairs  
(B) TYPE: nucleic acid

467607 "16029680

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

|            |            |            |            |            |            |     |
|------------|------------|------------|------------|------------|------------|-----|
| CGGCTCGAGC | TCTTAGGCTT | TGAAGCATTT | TTGINTGTGC | TCCCTGATCT | TCATGTCACC | 60  |
| ACCATGAAGT | TCTTAGCAGT | CCTGGTACTC | TTGGGAGTTT | CCATCTNTCT | GGTCTCTGCC | 120 |
| CAGAATCCGA | CAACAGCTGC | TNCAGCTGAC | ACGNATCCAG | CTACTGGTCC | TGCTGATGAT | 180 |
| GAAGCCCCTG | ANGCTGAAAC | CACTGCTGCT | GCNACCACTG | CGACCACTG  |            | 229 |

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 308 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

|            |            |            |            |            |            |     |
|------------|------------|------------|------------|------------|------------|-----|
| TAGGCTTTGA | AGCATTTTGT | TCTGTGCTCC | CTGATCTTCA | GGTCACCACC | ATGAAGTTCT | 60  |
| TAGCAGTCCT | GGTACTCTTG | GGAGTTTCCA | TCTTTCTGGT | CTCTGCCCAG | AATCCGACAA | 120 |
| CAGCTGCTCC | AGCTGACACG | TATCCAGCTA | CTGGTCCTGC | TGATGATGAA | GCCCCTGATG | 180 |
| CTGAAACCAC | TGCTGCTGCA | ACCACTGCGA | CCACTGCTGC | TCCTACCACT | GCAACCACCG | 240 |
| CTGCTTCTAC | CACTGCTCGT | AAAGACATTC | CAGTTTACC  | CAAATGGGTT | GGGGATCTTC | 300 |
| CGAATGGT   |            |            |            |            |            | 308 |

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 197 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

|            |            |            |            |            |            |     |
|------------|------------|------------|------------|------------|------------|-----|
| GTTTTACCCA | AATGGGTTGG | GGATCTCCCG | AATGGTAGAG | TGTGTCCCTG | AGATGGAATC | 60  |
| AGCTTGAGTC | TTCTGCAATT | GGTCACAAC  | ATTATGCTT  | CCTGTGATTT | CATCCAAC   | 120 |
| CTTACCTTGC | CTACGATATC | CCCTTTATCT | CTAATCAGTT | TATTTTCTTT | CAAATAAAAA | 180 |
| ATAACTATGA | GCAACAT    |            |            |            |            | 197 |

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 482 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: base\_polymorphism
- (B) LOCATION: 54
- (D) OTHER INFORMATION: /note= "'K' represents a G/T polymorphism at this position"

(ix) FEATURE:

- (A) NAME/KEY: base\_polymorphism
- (B) LOCATION: 312
- (D) OTHER INFORMATION: /note= "'Y' represents a C/T polymorphism at this position"

CCCTTTCTTT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

|            |            |            |             |            |            |     |
|------------|------------|------------|-------------|------------|------------|-----|
| CGGCTCGAGC | TCTTAGGCTT | TGAAGCATTT | TTGTCTGTGC  | TCCCTGATCT | TCAKGTCACC | 60  |
| ACCATGAAGT | TCTTAGCAGT | CCTGGTACTC | TTGGGAGTTT  | CCATCTTTCT | GGTCTCTGCC | 120 |
| CAGAATCCGA | CAACAGCTGC | TCCAGCTGAC | ACGTATCCAG  | CTACTGGTCC | TGCTGATGAT | 180 |
| GAAGCCCCTG | ATGCTGAAAC | CACTGCTGCT | GCAACCACTG  | CGACCACTGC | TGCTCCTACC | 240 |
| ACTGCAACCA | CCGCTGCTTC | TACCACTGCT | CGTAAAGACA  | TTCCAGTTTT | ACCCAAATGG | 300 |
| GTTGGGGATC | TYCCGAATGG | TAGAGTGTGT | CCCTGAGATG  | GAATCAGCTT | GAGTCTTCTG | 360 |
| CAATTGGTCA | CAACTATTCA | TGCTTCCTGT | GATTTTCATCC | AACTACTTAC | CTTGCCTACG | 420 |
| ATATCCCCTT | TATCTCTAAT | CAGTTTATTT | TCTTTCAAAT  | AAAAAATAAC | TATGAGCAAC | 480 |
| AT         |            |            |             |            |            | 482 |

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 553 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: base\_polymorphism
- (B) LOCATION: 543
- (D) OTHER INFORMATION: /note= "\R' represents an A/G polymorphism at this position"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

|            |            |             |            |            |             |     |
|------------|------------|-------------|------------|------------|-------------|-----|
| GAATTCGGCT | CGAGCGGCTC | GAGCTCTTAG  | GCTTTGAAGC | ATTTTGTCT  | GTGCTCCCTG  | 60  |
| ATCTTCATGT | CACCACCATG | AAGTTCTTAG  | CAGTCCTGGT | ACTCTTGGGA | GTTTCCATCT  | 120 |
| TTCTGGTCTC | TGCCCAGAAT | CCGACAACAG  | CTGCTCCAGC | TGACACGTAT | CCAGCTACTG  | 180 |
| GTCCTGCTGA | TGATGAAGCC | CCTGATGCTG  | AAACCACTGC | TGCTGCAACC | ACTGCGACCA  | 240 |
| CTGCTGCTCC | TACCACTGCA | ACCACCGCTG  | CTTCTACCAC | TGCTCGTAAA | GACATTCCAG  | 300 |
| TTTTACCCAA | ATGGGTTGGG | GATCTCCCGA  | ATGGTAGAGT | GTGTCCCTGA | GATGGAATCA  | 360 |
| GCTTGAGTCT | TCTGCAATTG | GTCACAACCTA | TTCATGCTTC | CTGTGATTTT | ATCCAACCTAC | 420 |
| TTACCTTGCC | TACGATATCC | CCTTTATCTC  | TAATCAGTTT | ATTTTCTTTT | AAATAAAAAA  | 480 |
| TAACTATGAG | CAACAAAAAA | AAAAAATAAC  | AAAAAATAAC | AAAAAATAAC | AAAAAATAAC  | 540 |
| AARGGGCGGC | CGC        |             |            |            |             | 553 |

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

|            |            |            |            |            |            |    |
|------------|------------|------------|------------|------------|------------|----|
| AGCTCGGAAT | TCCGAGCTTG | GATCCTCTAG | AGCGGCCGCC | GACTAGTGAG | CTCGTCGACC | 60 |
| CGGGAATT   |            |            |            |            |            | 68 |

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

|            |            |            |            |            |            |    |
|------------|------------|------------|------------|------------|------------|----|
| AATTAATTCC | CGGGTCGACG | AGCTCACTAG | TCGGCGGCCG | CTCTAGAGGA | TCCAAGCTCG | 60 |
|------------|------------|------------|------------|------------|------------|----|

GAATTCCG

68

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AGCGGATAAC AATTCACAC AGGA

24

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 18 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGTAAAACGA CGGCCAGT

18

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ACTGCTCGTA AAGACATTCC

20

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGGACACACT CTACCATTC

19

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AAGCCCCTGA TGCTGAAACC

20

ACTGCTCGTA AAGACATTCC

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGCAGAAGAC TCAAGCTGAT TCC

23

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CCCAGTCACG ACGTTGTAAA ACG

23

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GCGGCCGCCG GGACACACTC TACCATT

27

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 90 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Lys | Phe | Leu | Ala | Val | Leu | Val | Leu | Leu | Gly | Val | Ser | Ile | Phe | Leu |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Val | Ser | Ala | Gln | Asn | Pro | Thr | Thr | Ala | Ala | Pro | Ala | Asp | Thr | Tyr | Pro |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Ala | Thr | Gly | Pro | Ala | Asp | Asp | Glu | Ala | Pro | Asp | Ala | Glu | Thr | Thr | Ala |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |
| Ala | Ala | Thr | Thr | Ala | Thr | Thr | Ala | Ala | Pro | Thr | Thr | Ala | Thr | Thr | Ala |
|     |     | 50  |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |
| Ala | Ser | Thr | Thr | Ala | Arg | Lys | Asp | Ile | Pro | Val | Leu | Pro | Lys | Trp | Val |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Gly | Asp | Leu | Pro | Asn | Gly | Arg | Val | Cys | Pro |     |     |     |     |     |     |
|     |     |     |     | 85  |     |     |     | 90  |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO:17:



(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Gln | Asn | Pro | Thr | Thr | Ala | Ala | Pro | Ala | Asp | Thr | Tyr | Pro | Ala | Thr |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Gly | Pro | Ala | Asp | Asp | Glu | Ala | Pro | Asp | Ala | Glu | Thr | Thr | Ala | Ala | Ala |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Thr | Thr | Ala | Thr | Thr | Ala | Ala |     |     |     |     |     |     |     |     |     |
|     |     |     | 35  |     |     |     |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Thr | Ala | Thr | Thr | Ala | Ala | Pro | Thr | Thr | Ala | Thr | Thr | Ala | Ala | Ser |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Thr | Thr | Ala | Arg | Lys | Asp | Ile | Pro | Val | Leu | Pro | Lys | Trp | Val | Gly | Asp |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |
| Leu | Pro | Asn | Gly | Arg | Val | Cys |     |     |     |     |     |     |     |     |     |
|     |     |     | 35  |     |     |     |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Arg | Lys | Asp | Ile | Pro | Val | Leu | Pro | Lys | Trp | Val | Gly | Asp | Leu | Pro |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Asn | Gly | Arg | Val | Cys |     |     |     |     |     |     |     |     |     |     |     |
|     |     |     | 20  |     |     |     |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

46029680

Ala Ala Pro Ala Asp Thr Tyr Pro Ala Thr Gly Pro Ala Asp Asp Glu  
1 5 10 15  
Ala Pro Asp Ala Glu  
20

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Asp Tyr Lys Asp Asp Asp Asp Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn Met His Thr Glu His  
1 5 10 15  
His His His His  
20

We Claim:

1. A method of detecting the presence of a target BS106 polynucleotide  
5 in a test sample, comprising:
  - (a) contacting said test sample with at least one BS106-specific polynucleotide or complement thereof; and
  - (b) detecting the presence of said target BS106 polynucleotide in the test sample, wherein said BS106-specific polynucleotide has at least 50% identity to a  
10 polynucleotide selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.
2. The method of claim 1, wherein said target BS106 polynucleotide is  
15 attached to a solid phase prior to performing step (a).
3. A method for detecting mRNA of BS106 in a test sample, comprising:
  - (a) performing reverse transcription with at least one primer in order to  
20 produce cDNA;
  - (b) amplifying the cDNA obtained from step (a) using BS106 oligonucleotides as sense and antisense primers to obtain BS106 amplicon; and
  - (c) detecting the presence of said BS106 amplicon in the test sample, wherein the BS106 oligonucleotides utilized in steps (a) and (b) have at least 50%  
25 identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.
4. The method of claim 3, wherein said test sample is reacted with a  
30 solid phase prior to performing one of steps (a), (b), or (c).
5. The method of claim 3, wherein said detection step comprises utilizing a detectable label capable of generating a measurable signal.

6. A method of detecting a target BS106 polynucleotide in a test sample suspected of containing said target, comprising:

- (a) contacting said test sample with at least one BS106 oligonucleotide as a sense primer and with at least one BS106 oligonucleotide as an anti-sense primer  
5 and amplifying to obtain a first stage reaction product;
- (b) contacting said first stage reaction product with at least one other BS106 oligonucleotide to obtain a second stage reaction product, with the proviso that the other BS106 oligonucleotide is located 3' to the BS106 oligonucleotides utilized in step (a) and is complementary to said first stage reaction product; and  
10
- (c) detecting said second stage reaction product as an indication of the presence of the target BS106 polynucleotide, wherein the BS106 oligonucleotides utilized in steps (a) and (b) have at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or  
15 complements thereof.

7. The method of claim 6, wherein said test sample is reacted with a solid phase prior to performing one of steps (a), (b), or (c).

20 8. The method of claim 6, wherein said detection step comprises utilizing a detectable label capable of generating a measurable signal.

9. The method of claim 8, wherein said detectable label is reacted to a solid phase.  
25

10. A test kit useful for detecting BS106 polynucleotide in a test sample, comprising a container containing at least one BS106 polynucleotide having at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4,  
30 SEQUENCE ID NO 5, and fragments or complements thereof.

11. A purified polynucleotide or fragment thereof derived from a BS106 gene, wherein said polynucleotide is capable of selectively hybridizing to the nucleic acid of said BS106 gene and has at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID  
35

NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.

12. The purified polynucleotide of claim 11, wherein said polynucleotide  
5 is produced by recombinant techniques.

13. The purified polynucleotide of claim 11, wherein said polynucleotide is produced by synthetic techniques.

10 14. The purified polynucleotide of claim 11, wherein said polynucleotide comprises a sequence encoding at least one BS106 epitope.

15 15. A recombinant expression system comprising a nucleic acid sequence that includes an open reading frame derived from BS106 operably linked to a control sequence compatible with a desired host, wherein said nucleic acid sequence has at least 50% identity to a sequence selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.

20 16. A cell transfected with the recombinant expression system of claim 15.

25 17. A BS106 polypeptide having at least 50% identity with an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments of SEQUENCE ID NOS 17-20.

30 18. The polypeptide of claim 17, wherein said polypeptide is produced by recombinant techniques.

19. The polypeptide of claim 17, wherein said polypeptide is produced by synthetic techniques.

35 20. An antibody which specifically binds to at least one BS106 epitope, wherein said BS106 epitope is derived from an amino acid sequence having at least

2025 RELEASE UNDER E.O. 14176

50% identity to an amino acid sequence selected from the group consisting of  
SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18,  
SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

5           21.     An assay kit for determining the presence of BS106 antigen or anti-  
BS106 antibody in a test sample, comprising a container containing a BS106  
polypeptide having at least 50% identity with an amino acid sequence selected from  
the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17,  
SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and  
10    fragments thereof.

          22.     The assay kit of claim 21, wherein said polypeptide is attached to a  
solid phase.

15           23.     An assay kit for determining the presence of BS106 antigen in a test  
sample, comprising a container containing an antibody which specifically binds to a  
BS106 antigen which comprises at least one BS106 epitope.

          24.     The kit of claim 23, wherein said antibody is attached to a solid  
20    phase.

          25.     A method for producing a polypeptide comprising at least one BS106  
epitope, said method comprising incubating host cells that have been transfected with  
an expression vector containing a polynucleotide sequence encoding a polypeptide,  
25    wherein said polypeptide comprises an amino acid sequence having at least 50%  
identity to an amino acid sequence selected from the group consisting of  
SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18,  
SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

30           26.     A method for detecting BS106 antigen in a test sample suspected of  
containing said BS106 antigen, comprising:

          (a)     contacting the test sample with an antibody or fragment thereof which  
specifically binds to at least one epitope of a BS106 antigen selected from the group  
consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO  
35    18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof,

“SECRET” 46029680

wherein said contacting is carried out for a time and under conditions sufficient for the formation of antibody/antigen complexes; and

(b) detecting the presence of said complexes as an indication of the presence of said BS106 antigen.

5

27. The method of claim 26, wherein said antibody is attached to a solid phase.

28. A method for detecting the presence of antibodies specific for a BS106 antigen in a test sample suspected of containing such antibodies, said method comprising:

(a) contacting the test sample with a BS106 polypeptide, wherein said BS106 polypeptide contains at least one BS106 epitope derived from an amino acid sequence or fragment thereof having at least 50% identity to an amino acid sequence selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof, and further wherein said contacting is carried out for a time and under conditions sufficient to allow antigen/antibody complexes to form; and

(b) detecting said complexes.

20

29. The method of claim 28, wherein said BS106 polypeptide is attached to a solid phase.

30. A cell transfected with a nucleic acid sequence encoding at least one BS106 epitope, wherein said nucleic acid sequence is selected from the group consisting of SEQUENCE ID NO 1, SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE ID NO 5, and fragments or complements thereof.

31. A method for producing antibodies which specifically bind to BS106 antigen, comprising administering to an individual an isolated immunogenic polypeptide or fragment thereof in an amount sufficient to elicit an immune response, wherein said immunogenic polypeptide comprises at least one BS106 epitope and has at least 50% identity to a sequence selected from the group consisting of

46 FEB " 16029680

SEQUENCE ID NO 16, SEQUENCE ID NO 17, SEQUENCE ID NO 18,  
SEQUENCE ID NO 19, SEQUENCE ID NO 20, and fragments thereof.

32. A method for producing antibodies which specifically bind to BS106  
5 antigen, comprising administering to a mammal a plasmid comprising a sequence  
which encodes at least one BS106 epitope derived from a polypeptide having an  
amino acid sequence selected from the group consisting of SEQUENCE ID NO 16,  
SEQUENCE ID NO 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19,  
SEQUENCE ID NO 20, and fragments thereof.

10

33. A composition of matter comprising a BS106 polynucleotide or  
fragment thereof, wherein said polynucleotide has at least 50% identity to a  
polynucleotide selected from the group consisting of SEQUENCE ID NO 1,  
SEQUENCE ID NO 2, SEQUENCE ID NO 3, SEQUENCE ID NO 4, SEQUENCE  
15 ID NO 5, and fragments or complements thereof.

34. A composition of matter comprising a polypeptide containing at least  
one BS106 epitope, wherein said polypeptide has at least 50% identity to a sequence  
selected from the group consisting of SEQUENCE ID NO 16, SEQUENCE ID NO  
20 17, SEQUENCE ID NO 18, SEQUENCE ID NO 19, SEQUENCE ID NO 20, and  
fragments of SEQUENCE ID NOS 17-20.

35. The test kit of claim 10 further comprising a container with tools  
useful for collection of said sample, wherein the tools are selected from the group  
25 consisting of lancets, absorbent paper, cloth, swabs and cups.

36. The assay kit of claim 21 further comprising a container with tools  
useful for collection of said sample, wherein the tools are selected from the group  
consisting of lancets, absorbent paper, cloth, swabs and cups.

30

37. The test kit of claim 23 further comprising a container with tools  
useful for collection of said sample, wherein the tools are selected from the group  
consisting of lancets, absorbent paper, cloth, swabs and cups.

08992994 16029680



38. A gene or fragment thereof which codes for a BS106 protein which comprises an amino acid sequence with at least 50% identity with SEQUENCE ID NO 16.

5 39. A gene or fragment thereof comprising DNA having at least 50% identity with SEQUENCE ID NO 4 or SEQUENCE ID NO 5.

467507-4623680

REAGENTS AND METHODS USEFUL FOR  
DETECTING DISEASES OF THE BREAST

5

Abstract of the Disclosure

10

15 A set of contiguous and partially overlapping RNA sequences and polypeptides encoded thereby, designated as BS106 and transcribed from breast tissue is described. These sequences are useful for the detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, or determining the predisposition of an individual to diseases and conditions of the breast such as breast cancer. Also provided are antibodies which specifically bind to BS106-encoded polypeptide or protein, and agonists or inhibitors which prevent action of the tissue-specific BS106 polypeptide, which molecules are useful for the therapeutic treatment of breast diseases, tumors or metastases.

46TET-4623680

# FIGURE 1

|           |            |            |            |             |             |
|-----------|------------|------------|------------|-------------|-------------|
| >1662885  | CGGCTCGAGC | TCTTAGGCTT | TGAAGCATTT | TTGTNTGTGC  | TCCCTGATCT  |
| >893988   |            | TAGGCTT    | TGAAGCATTT | TTGTCTGTGC  | TCCCTGATCT  |
| Consensus | CGGCTCGAGC | TCTTAGGCTT | TGAAGCATTT | TTGTCTGTGC  | TCCCTGATCT  |
|           |            |            |            |             |             |
| >1662885  | TCATGTCACC | ACCATGAAGT | TCTTAGCAGT | CCTGGTACTC  | TTGGGAGTTT  |
| >893988   | TCAGGTCACC | ACCATGAAGT | TCTTAGCAGT | CCTGGTACTC  | TTGGGAGTTT  |
| Consensus | TCAGGTCACC | ACCATGAAGT | TCTTAGCAGT | CCTGGTACTC  | TTGGGAGTTT  |
|           |            |            |            |             |             |
| >1662885  | CCATCTNTCT | GGTCTCTGCC | CAGAATCCGA | CAACAGCTGC  | TNCAGCTGAC  |
| >893988   | CCATCTTTCT | GGTCTCTGCC | CAGAATCCGA | CAACAGCTGC  | TCCAGCTGAC  |
| Consensus | CCATCTTTCT | GGTCTCTGCC | CAGAATCCGA | CAACAGCTGC  | TCCAGCTGAC  |
|           |            |            |            |             |             |
| >1662885  | ACGNATCCAG | CTACTGGTCC | TGCTGATGAT | GAAGCCCCCTG | ANGCTGAAAC  |
| >893988   | ACGTATCCAG | CTACTGGTCC | TGCTGATGAT | GAAGCCCCCTG | ATGCTGAAAC  |
| Consensus | ACGTATCCAG | CTACTGGTCC | TGCTGATGAT | GAAGCCCCCTG | ATGCTGAAAC  |
|           |            |            |            |             |             |
| >1662885  | CACTGCTGCT | GCNACCACTG | CGACCACTG  |             |             |
| >893988   | CACTGCTGCT | GCAACCACTG | CGACCACTGC | TGCTCCTACC  | ACTGCAACCA  |
| Consensus | CACTGCTGCT | GCAACCACTG | CGACCACTGC | TGCTCCTACC  | ACTGCAACCA  |
|           |            |            |            |             |             |
| >893988   | CCGCTGCTTC | TACCACTGCT | CGTAAAGACA | TTCCAGTTTT  | ACCCAAATGG  |
| >1209814  |            |            |            | GTTTT       | ACCCAAATGG  |
| Consensus | CCGCTGCTTC | TACCACTGCT | CGTAAAGACA | TTCCAGTTTT  | ACCCAAATGG  |
|           |            |            |            |             |             |
| >893988   | GTTGGGGATC | TTCCGAATGG | T          |             |             |
| >1209814  | GTTGGGGATC | TCCCGAATGG | TAGAGTGTGT | CCCTGAGATG  | GAATCAGCTT  |
| Consensus | GTTGGGGATC | TYCCGAATGG | TAGAGTGTGT | CCCTGAGATG  | GAATCAGCTT  |
|           |            |            |            |             |             |
| >1209814  | GAGTCTTCTG | CAATTGGTCA | CAACTATTCA | TGCTTCCTGT  | GATTTTCATCC |
| Consensus | GAGTCTTCTG | CAATTGGTCA | CAACTATTCA | TGCTTCCTGT  | GATTTTCATCC |
|           |            |            |            |             |             |
| >1209814  | AACTACTTAC | CTTGCCTACG | ATATCCCCTT | TATCTCTAAT  | CAGTTTATTT  |
| Consensus | AACTACTTAC | CTTGCCTACG | ATATCCCCTT | TATCTCTAAT  | CAGTTTATTT  |
|           |            |            |            |             |             |
| >1209814  | TCTTTCAAAT | AAAAAATAAC | TATGAGCAAC | AT          |             |
| Consensus | TCTTTCAAAT | AAAAAATAAC | TATGAGCAAC | AT          |             |

FIGURE 2




Figure 3A





FIGURE 3B



| Lane | Tissue        |
|------|---------------|
| 1    | Normal Breast |
| 2    | Normal Breast |
| 3    | Normal Breast |
| 4    | Normal Breast |
| 5    | Normal Breast |
| 6    | Normal Breast |
| 7    | Cancer Breast |
| 8    | Cancer Breast |
| 9    | Cancer Breast |
| 10   | Cancer Breast |
| 11   | Cancer Breast |
| 12   | Cancer Breast |

**Figure 4A**





Figure 4B





COMBINED DECLARATION AND POWER OF ATTORNEY  
FOR CONTINUATION-IN-PART APPLICATION

AS A BELOW-NAMED INVENTOR, I HEREBY DECLARE THAT:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: REAGENTS AND METHODS USEFUL FOR DETECTING DISEASES OF THE BREAST, the specification of which

  X   is attached hereto  
       was filed on

and assigned Serial No.    and was amended on   .

I HAVE REVIEWED AND UNDERSTAND THE CONTENTS OF THE ABOVE-IDENTIFIED SPECIFICATION, INCLUDING THE CLAIMS, AS AMENDED BY ANY AMENDMENT REFERRED TO ABOVE.

I acknowledge and understand that I am an individual who has a duty to disclose information which is material to the patentability of the claims of this application in accordance with Title 37, Code of Federal Regulations, §§ 1.56(a) and (b) which state:

"(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is cancelled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

(1) prior art cited in search reports of a foreign patent office in a counterpart application, and

(2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

(1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim; or

(2) It refutes, or is inconsistent with, a position the applicant takes in:

(i) Opposing an argument of unpatentability relied on by the Office,

or

(ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability."

I do not know and do not believe this invention was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to said application. This invention was not in public use or on sale in the United States of America more than one year prior to this application. This invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on any application filed by me or my legal representatives or assigns more than six months prior to this application.

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) and (b) set forth above which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.: 08/742,067

Filing Date: October 31, 1996

Status (patented, pending, abandoned): pending

Application Serial No.:

Filing Date:

Status (patented, pending, abandoned):

As to the subject matter of this application which is common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to said earlier application, or in public use or on sale in the United States of America more than one year prior to said earlier application; that said common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to said earlier application; and that the earliest application(s) for patent or inventor's certificate on said invention filed by me or my legal representatives or assigns in any country foreign to the United States of America is identified below, as well as all other such applications (if any) filed more than twelve months prior to the filing date of this application:

None.

The priority of the earliest application(s) (if any) filed within a year prior to said pending prior application is hereby claimed under 35 U.S.C. § 119.

As to the subject matter of this application which is not common to said earlier application, I do not know and do not believe that the same was ever known or used in the United States of America before my or our invention thereof or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to the date of this application, or in public use or on sale in the United States of America more than one year prior to the date of this application, and that said subject matter has not been patented or made the subject of an inventor's certificate issued in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to the date of this application, and that the earliest application(s) for patent or inventor's certificate on said subject matter filed by me or my legal representatives or assigns in any country foreign to the United States of America is identified below, as well as all other such application(s) (if any) filed more than twelve months prior to the filing date of this application:

None.

The priority of the earliest application(s) (if any) filed within a year to this application is hereby claimed under 35 U.S.C. § 119.

I hereby appoint the following attorneys and agents to prosecute that application and to transact all business in the Patent and Trademark Office connected therewith and to file, to prosecute and to transact all business in connection with all patent applications directed to the invention:

Mona Anand, Reg. No. 34,537  
Mark C. Bach, Reg. No. 34,766  
Cheryl L. Becker, Reg. No. 35,441  
Thomas D. Brainard, Reg. No. 32,459  
Thomas M. Breining, Reg. No. 29,897  
Monte R. Browder, Reg. No. 36,761  
Dianne Casuto, Reg. No. P-40,943  
Steven R. Crowley, Reg. No. 31,604  
Andreas M. Danckers, Reg. No. 32,652  
Richard A. Elder, Reg. No. 30,255  
Roberta L. Robins, Reg. No. 33,208

Matthew R. Hooper, Reg. No. 31,108  
James D. McNeil, Reg. No. 26,204  
Lawrence S. Pope, Reg. No. 26,791  
Nicholas A. Poulos, Reg. No. 30,209  
Priscilla E. Porembski, Reg. No. 33,207  
Gregory W. Steele, Reg. No. 33,796  
Michael J. Ward, Reg. No. 37,960  
David L. Weinstein, Reg. No. 28,128  
Brian L. Woodworth, Reg. No. 33,137  
Paul D. Yasger, Reg. No. 37,477  
Thomas P. McCracken, Reg. No. 38,548

Send Correspondence to:

Steven F. Weinstock  
Abbott Laboratories  
D-377/AP6D  
100 Abbott Park Road  
Abbott Park, Illinois 60064-3500

Direct telephone calls to:

Cheryl L. Becker at 847-937-6100

Name (first, middle, last): Patricia A. BILLING-MEDEL

Post Office Address: 1427 Sherwood Court, Gurnee, IL 60031

Residence: Gurnee, IL 60031

Citizenship: US

Name (first, middle, last): Maurice COHEN

Post Office Address: 2026 Deerfield Road, Highland Park, IL 60035

Residence: Highland Park, IL 60035

Citizenship: US

Name (first, middle, last): Tracey L. COLPITTS

Post Office Address: 34365 North Circle Drive, Round Lake, IL 60073

Residence: Round Lake, IL 60073

Citizenship: Canada

Name (first, middle, last): Paula N. FRIEDMAN

Post Office Address: 462 Cumnor Court, Deerfield, IL 60015

Residence: Deerfield, IL 60015

Citizenship: US

Name (first, middle, last): Julian GORDON

Post Office Address: 307 East Sheridan Road, Lake Bluff, IL 60044

Residence: Lake Bluff, IL 60044

Citizenship: US

Name (first, middle, last): Edward N. GRANADOS

Post Office Address: 19 Montgomery Lane, Vernon Hills, IL 60061

Residence: Vernon Hills, IL 60061

Citizenship: US

Name (first, middle, last): Steven C. HODGES

Post Office Address: 169 Stonegate Road, Buffalo Grove, IL 60089

Residence: Buffalo Grove, IL 60089

Citizenship: US

Name (first, middle, last): Michael R. KLASS

Post Office Address: 1606 Mulberry Drive, Libertyville, IL 60048

Residence: Libertyville, IL 60048

Citizenship: US

Name (first, middle, last): Jon D. KRATOCHVIL

Post Office Address: 7101 Fifth Avenue, Kenosha, WI 53143

Residence: Kenosha, WI 53143

Citizenship: US

Name (first, middle, last): Lisa ROBERTS-RAPP

Post Office Address: 2090 Westfield Drive, Gurnee, IL 60031

Residence: Gurnee, IL 60031

Citizenship: US

Name (first, middle, last): John C. RUSSELL

Post Office Address: 8275 64th Court, Kenosha, WI 53142

Residence: Kenosha, WI 53142

Citizenship: US

Name (first, middle, last): Steven D. STROUPE

Post Office Address: 945 Wilshire Drive, Libertyville, IL 60048

Residence: Libertyville, IL 60048

Citizenship: US

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that all statements made herein were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States

Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Patricia A. BILLING-MEDEL

Maurice COHEN

Tracey L. COLPITTS

Paula N. FRIEDMAN

Julian GORDON

Edward N. GRANADOS

Steven C. HODGES

Michael R. KLASS

Jon D. KRATOCHVIL

Lisa ROBERTS-RAPP

John C. RUSSELL

Steven D. STROUPE

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED